

Jack polynomials in superspace

Patrick Desrosiers*

Département de Physique, de Génie Physique et d'Optique,
Université Laval,
Québec, Canada, G1K 7P4.

Luc Lapointe†

Instituto de Matemática y Física,
Universidad de Talca,
Casilla 747, Talca, Chile.

Pierre Mathieu‡

Département de Physique, de Génie Physique et d'Optique,
Université Laval,
Québec, Canada, G1K 7P4.

October 2002

Abstract

This work initiates the study of *orthogonal* symmetric polynomials in superspace. Here we present two approaches leading to a family of orthogonal polynomials in superspace that generalize the Jack polynomials. The first approach relies on previous work by the authors in which eigenfunctions of the supersymmetric extension of the trigonometric Calogero-Moser-Sutherland Hamiltonian were constructed. Orthogonal eigenfunctions are now obtained by diagonalizing the first nontrivial element of a bosonic tower of commuting conserved charges not containing this Hamiltonian. Quite remarkably, the expansion coefficients of these orthogonal eigenfunctions in the supermonomial basis are stable with respect to the number of variables. The second and more direct approach amounts to symmetrize products of non-symmetric Jack polynomials with monomials in the fermionic variables. This time, the orthogonality is inherited from the orthogonality of the non-symmetric Jack polynomials, and the value of the norm is given explicitly.

*pdesrosi@phy.ulaval.ca

†lapointe@inst-mat.otalca.cl

‡pmathieu@phy.ulaval.ca

1 Introduction

A natural direction in which the theory of orthogonal symmetric polynomials can be generalized is to consider its extension to superspace. One possible approach for such an extension is to consider polynomials involving fermionic (i.e., Grassmannian) variables, or superpolynomials, that arise from physically relevant eigenvalue problems invariant under supersymmetry.

In many respects (physical and mathematical), one of the most fundamental bases of symmetric orthogonal polynomials is that of the Jack polynomials. This work is concerned with their orthogonality-preserving extension to superspace.

A basic requirement of Jack superpolynomials is that they reduce to Jack polynomials when the fermionic variables are set to zero. Another requirement is that they be solutions of the supersymmetric generalization of the eigenvalue problem characterizing the Jack polynomials. More precisely, Jack polynomials are eigenfunctions of the trigonometric Calogero-Moser-Sutherland (tCMS) model (see e.g., [1] and [2, 3] for properties of the Jack polynomials). Jack superpolynomials must thus be eigenfunctions of the supersymmetric extension of the trigonometric Calogero-Moser-Sutherland (stCMS) model [4].¹

This eigenfunction characterization, as in the non-fermionic case, does not uniquely define Jack superpolynomials. A triangular decomposition, in terms of a superspace extension of the symmetric monomial functions (or supermonomials), must be imposed. Unique eigenfunctions, $\mathcal{J}_\Lambda$, defined according to such a triangular decomposition, were constructed in [4, 5], and called Jack superpolynomials. They are indexed by superpartitions $\Lambda = (\Lambda^a; \Lambda^s)$, composed of a partition Λ^a with distinct parts, and a usual partition Λ^s . The number of entries in Λ^a characterizes the fermion sector (i.e., the number of anticommuting variables appearing in every term of the expansion of $\mathcal{J}_\Lambda$).

The integrability of the tCMS model also makes Jack polynomials eigenfunctions of a family of N independent commuting quantities, where N is the number of variables. We prove in this article that, similarly, Jack superpolynomials are eigenfunctions of a whole tower of commuting conserved charges, denoted H_n , for $n = 1, \dots, N$, where H_2 is the Hamiltonian of the stCMS model. The proof relies heavily on the remarkable fact that, if we consider the restriction to the space of superpolynomials symmetric under the simultaneous interchange of any pair of bosonic and fermionic variables, these charges can be expressed using Dunkl operators as $\mathcal{H}_n = \sum_{i=1}^N (\mathcal{D}_i)^n$. That is, under this restriction, H_n is equivalent to $\mathcal{H}_n$.

Now, even though Jack superpolynomials are eigenfunctions of N commuting conserved charges, degeneracies are still present. Indeed, two distinct superpolynomials labeled by two different superpartitions built out of the same set of N integers (but distributed differently among the two partitions Λ^a and Λ^s) have identical $\mathcal{H}_n$ eigenvalues. As a result, the Jack superpolynomials $\mathcal{J}_\Lambda$ of [4, 5] are *not orthogonal* under the scalar product (19) with respect to which $\mathcal{H}_n$ is self-adjoint.

The Gram-Schmidt orthogonalization procedure can of course be used to construct orthogonal superpolynomials, but a general pattern is not likely to appear using this construction. The question is thus whether we can naturally define a family of orthogonal superpolynomials. The answer to this question lies in the following observation. By extending the usual tCMS model with N bosonic degrees of freedom to the supersymmetric case, we have introduced N new degrees of freedom. Integrability leads in this case to the appearance of new conserved charges. Indeed, there are $3N$ new conserved charges, new in the sense that they all disappear when the fermionic variables vanish [4]. Among these new charges, $2N$ of them are fermionic, that is, they change the fermion number of the function on which they act. However, the remaining N charges are bosonic, mutually commute, and do not affect the fermion number. These charges, denoted I_n , $n = 1, \dots, N$, are thus natural candidates for extra operators that may lift the degeneracy of the Jack superpolynomials, and thereby produce orthogonal combinations of these superpolynomials.

This expectation indeed materializes. Actually, to construct orthogonal superpolynomials it suffices to consider the action of the charge I_1 , or equivalently, its Dunkl-operator version $\mathcal{I}_1$ in the space of symmetric

¹An extensive list of references on the CMS model and its supersymmetric extension can be found in [4].

superpolynomials. Its action is also triangular, but with respect to an ordering on superpartitions stronger than the one introduced in [5]. Knowing the action of $\mathcal{I}_1$ explicitly on Jack superpolynomials allows to define orthogonal fermionic extensions, J_Λ , of the usual Jack polynomials. Moreover, it also leads to determinantal formulas for the expansion coefficients of the orthogonal Jack superpolynomials J_Λ in terms of Jack superpolynomials $\mathcal{J}_\Omega$.

The program that we just sketched is the subject of the first part of this paper (up to section 8). It is in line with our previous work [4, 5], and can be viewed as its natural completion. We stress that it is also very explicit in that the precise relation between the orthogonal Jack superpolynomials J_Λ with the old $\mathcal{J}_\Lambda$ is provided, and that closed form expressions for the latter, in the supermonomial basis, were already obtained in [5]. The construction is also ‘physical’: the quantum many-body problem and its underlying integrability structure is the guiding tool used to identify a complete set of simultaneously diagonalizable operators.

This leads us to our first characterization of the orthogonal Jack superpolynomials:

Theorem 1. (See Theorems 22 and 31). *The orthogonal Jack superpolynomial J_Λ is the unique function satisfying:*

$$\mathcal{H}_2 J_\Lambda = \varepsilon_\Lambda J_\Lambda, \quad \mathcal{I}_1 J_\Lambda = \epsilon_\Lambda J_\Lambda, \quad \text{and} \quad J_\Lambda = m_\Lambda + \sum_{\Omega < \Lambda} c_{\Lambda\Omega}(\beta) m_\Omega, \quad (1)$$

where ε_Λ and ϵ_Λ are defined in Lemma 20 and Theorem 28 respectively, while the ordering on superpartitions is introduced in Definition 7.

In the second part of the paper (which is essentially section 9) we propose a much more direct, although less explicit, construction of the orthogonal Jack superpolynomials. The starting point is not anymore the extension of the usual Jack polynomial eigenvalue problem, but rather a symmetrization process performed on the non-symmetric Jack polynomials [6, 7], suitably dressed with products of fermionic variables.

Recall that the non-symmetric Jack polynomials E_λ (where λ is now a composition) are eigenfunctions of the $\mathcal{D}_i$ operators [6], and, from the self-adjointness of these operators, orthogonal. On the other hand, as already pointed out, the stCMS commuting conserved charges can all be expressed in term of the $\mathcal{D}_i$ ’s. This naturally suggests a very direct path for the construction of the common eigenfunctions of all the commuting stCMS charges: start with E_λ , add a fermionic-monomial prefactor and symmetrize with respect to both types of variables. Quite remarkably, this indeed produces the orthogonal Jack superpolynomials J_Λ (with Λ determined by λ and the fermionic number). The advantage of this construction is that orthogonality is built in, and preserved by the symmetrization.

This second construction (especially the argument in the proof of Theorem 41) leads to another characterization of the orthogonal Jack superpolynomials:

Theorem 2. (See Theorems 35 and 41). *The orthogonal Jack superpolynomials are the unique functions satisfying:*

$$\langle J_\Lambda, J_\Omega \rangle_\beta \propto \delta_{\Lambda\Omega}, \quad \text{and} \quad J_\Lambda = m_\Lambda + \sum_{\Omega < \Lambda} c_{\Lambda\Omega}(\beta) m_\Omega. \quad (2)$$

Our two characterizations of the orthogonal Jack superpolynomials thus extend the two most common definitions of Jack polynomials.

An important clarification is in order concerning the second construction. Because of the anticommuting nature of the fermionic variables, a symmetrized superpolynomial that contains m fermionic variables is necessarily an antisymmetric function of the m corresponding bosonic variables, in addition to be symmetric in the remaining bosonic variables. In other words, viewed solely as functions of the bosonic variables, the symmetric superpolynomials can be decomposed as a sum over polynomials with mixed symmetry properties such as those studied in [8, 9], but with coefficients involving fermionic variables. This, however, does not mean that the theory of symmetric superpolynomials is only a special case of the theory of polynomials with mixed symmetry. Indeed, by symmetrizing over all the variables, including the fermionic ones, the

antisymmetrized bosonic variables will be dependent of the particular term under consideration, ensuring that the net result is a brand new object. Let us illustrate this comment by comparing a four-variable monomial function with mixed symmetry with its supermonomial counterpart. Take the four variables to be x_1, x_2 , and y_1, y_2 , and let the x_i variables be symmetrized while the y_i are antisymmetrized. Let also the x and y parts of the monomial be parametrized by the partitions $(2, 1)$ and $(3, 1)$ respectively. The associated monomial with mixed symmetry reads:

$$\tilde{m}_{(3,1)_a, (2,1)_s} = (y_1^3 y_2 - y_1 y_2^3)(x_1^2 x_2 + x_1 x_2^2) \quad (3)$$

The corresponding supermonomial has four bosonic and four fermionic variables denoted z_i and θ_i respectively, with $i = 1, \dots, 4$ and $\theta_i \theta_j = -\theta_j \theta_i$. Given the superpartition $(3, 1; 2, 1)$, it reads as

$$\begin{aligned} m_{(3,1;2,1)} = & \theta_1 \theta_2 (z_1^3 z_2 - z_1 z_2^3)(z_3^2 z_4 + z_3 z_4^2) + \theta_1 \theta_3 (z_1^3 z_3 - z_1 z_3^3)(z_2^2 z_4 + z_2 z_4^2) \\ & + \theta_1 \theta_4 (z_1^3 z_4 - z_1 z_4^3)(z_2^2 z_3 + z_2 z_3^2) + \theta_2 \theta_3 (z_2^3 z_3 - z_2 z_3^3)(z_1^2 z_4 + z_1 z_4^2) \\ & + \theta_2 \theta_4 (z_2^3 z_4 - z_2 z_4^3)(z_1^2 z_3 + z_1 z_3^2) + \theta_3 \theta_4 (z_3^3 z_4 - z_3 z_4^3)(z_1^2 z_2 + z_1 z_2^2) \end{aligned} \quad (4)$$

Clearly, each bosonic component of this expression corresponds to a monomial with mixed symmetry of type (3). The main point is that the supermonomial is the sum of all these mixed symmetry monomials, each with its appropriate fermionic-monomial prefactor. These prefactors have drastic effects say, when multiplying supermonomials together, due to the fermionic nature of their constituents. The multiplication of polynomials reveals in a rather critical way one aspect of the difference between the polynomials with mixed symmetry and superpolynomials: the product of two polynomials of the former type cannot be decomposed into a linear combination of polynomials with mixed symmetry, that is, there is no ring structure. This is not so for the superpolynomials. It should thus be crystal clear that Jack superpolynomials are not simply Jack polynomials with mixed symmetry properties in disguised form. However, pinpointing the relationship between these two types of objects is technically important since it allows to use the results of [8] on the norm of the Jack polynomials with mixed symmetry to obtain, in a rather direct way, the norm of the orthogonal Jack superpolynomials.

The presentation of the two approaches is preceded by four sections in which we introduce our notation, define our basic superobjects and derive relevant properties. Section 6 and 7 deal respectively with the construction of the $\mathcal{H}_n$ and $\mathcal{I}_n$ eigenfunctions. The common eigenfunctions are shown to be orthogonal in Section 8. The alternative construction based on the non-symmetric Jack polynomials is the subject of Section 9.

In Appendix A, we present a number of examples of Jack superpolynomials, including a detailed computation based on the determinantal formula. These examples illustrate a nice property of the orthogonal Jack superpolynomials: they do not depend upon N (when N is sufficiently large). In other words, their expansion coefficients in the supermonomial basis are independent of the number of variables. This property can of course be obtained from the explicit formulas, but it is not at once manifest.

Finally, various natural extensions of this work are mentioned in the conclusion.

Note that for the readers not particularly interested in the ‘physical’ construction relying on the structure of the integrable supersymmetric stCMS model and its conserved charges, reading Section 2, and the first two definitions of Section 4 is sufficient to understand Section 9.

2 Basic definitions

For $i, j \in \{1, \dots, N\}$, let K_{ij} be the operator that exchanges the variables z_i and z_j . Similarly, let κ_{ij} exchange the anticommuting variables θ_i and θ_j . Their action on a superfunction $f(z, \theta)$ is thus

$$\begin{aligned} K_{ij} f(\dots, z_i, \dots, z_j, \dots, \theta_i, \dots, \theta_j, \dots) &= f(\dots, z_j, \dots, z_i, \dots, \theta_i, \dots, \theta_j, \dots) , \\ \kappa_{ij} f(\dots, z_i, \dots, z_j, \dots, \theta_i, \dots, \theta_j, \dots) &= f(\dots, z_i, \dots, z_j, \dots, \theta_j, \dots, \theta_i, \dots) . \end{aligned} \quad (5)$$

Each of these sets of operators generates a realization of the permutation group S_N . Since the K_{ij} 's and the κ_{ij} 's commute, the operators $\mathcal{K}_{ij} = K_{ij}\kappa_{ij}$, acting as

$$\mathcal{K}_{ij}f(\dots, z_i, \dots, z_j, \dots, \theta_i, \dots, \theta_j, \dots) = f(\dots, z_j, \dots, z_i, \dots, \theta_j, \dots, \theta_i, \dots), \quad (6)$$

are also seen to generate a realization of S_N .

Let P be the space of polynomials in the variables $\theta_1, \dots, \theta_N$ and $z_1, \dots, z_N$. We will denote by P^{S_N} the subspace of P of polynomials invariant under the simultaneous exchange of any pair of variables $\theta_i \leftrightarrow \theta_j$ and $z_i \leftrightarrow z_j$. A polynomial $f \in P$ thus belongs to P^{S_N} iff $\mathcal{K}_\sigma f = f$ for any $\sigma \in S_N$.²

Let $I = \{i_1, \dots, i_m\}$, ($1 \leq i_1 < i_2 < \dots < i_m \leq N$) be an ordered set of integers, and let λ be a composition with N parts, that is, a sequence of N nonnegative integers (*e.g.*, if $N = 5$, one possible composition is (20134))³. A natural basis of P is provided by the monomials $\theta_I z^\lambda$, where

$$\theta_I = \theta_{\{i_1, \dots, i_m\}} = \theta_{i_1} \cdots \theta_{i_m}, \quad z^\lambda = z_1^{\lambda_1} \cdots z_N^{\lambda_N}. \quad (7)$$

If I has m entries, $\theta_I z^\lambda$ is said to belong to the m -fermion sector.

A superpartition Λ in the m -fermion sector is made of a partition Λ^a whose parts are all distinct, and of a usual partition Λ^s , that is,

$$\Lambda = (\Lambda_1, \dots, \Lambda_m; \Lambda_{m+1}, \dots, \Lambda_N) = (\Lambda^a; \Lambda^s), \quad (8)$$

with

$$\Lambda^a = (\Lambda_1, \dots, \Lambda_m), \quad \Lambda_i > \Lambda_{i+1} \geq 0, \quad i = 1, \dots, m-1, \quad (9)$$

and

$$\Lambda^s = (\Lambda_{m+1}, \dots, \Lambda_N), \quad \Lambda_i \geq \Lambda_{i+1} \geq 0, \quad i = m+1, \dots, N-1. \quad (10)$$

In the zero-fermion sector, the semicolon is omitted and Λ reduces to Λ^s . We often write the degree of a superpartition as $n = |\Lambda| = \sum_{i=1}^N \Lambda_i$. Given a supercomposition $\gamma = (\gamma_a; \gamma_s) = (\gamma_1, \dots, \gamma_m; \gamma_{m+1}, \dots, \gamma_N)$, we will denote by $\bar{\gamma}$ the superpartition whose antisymmetric part is the rearrangement of $(\gamma_1, \dots, \gamma_m)$ and whose symmetric part is the rearrangement of $(\gamma_{m+1}, \dots, \gamma_N)$. Denoting by λ^+ the partition obtained by the rearrangement of the entries of any composition λ , we have

$$\overline{(\gamma_a; \gamma_s)} = (\gamma_a^+; \gamma_s^+), \quad (11)$$

which we can illustrate with the example:

$$\overline{(1, 4, 2; 2, 5, 1, 3)} = (4, 2, 1; 5, 3, 2, 1). \quad (12)$$

Furthermore, σ_γ will stand for the element of S_N that sends γ to $\bar{\gamma}$, that is $\sigma_\gamma \gamma = \bar{\gamma}$. Note that we can always choose σ_γ such that $\sigma_\gamma = \sigma_\gamma^a \sigma_\gamma^s$, with σ_γ^a and σ_γ^s permutations of $\{1, \dots, m\}$ and $\{m+1, \dots, N\}$ respectively.

If we delete the semi-colon in a superpartition Λ , we obtain an ordinary composition that we will denote as Λ_c :

$$\Lambda = (\Lambda_1, \dots, \Lambda_m; \Lambda_{m+1}, \dots, \Lambda_N) \implies \Lambda_c = (\Lambda_1, \dots, \Lambda_N). \quad (13)$$

²It is understood that the product decomposition of $\mathcal{K}_\sigma$ into elementary permutations $\mathcal{K}_{i,i+1}$ follows the decomposition of σ into elementary transpositions $\sigma_i = (i, i+1)$. In other words, $\mathcal{K}_\sigma$ is a realization on superspace variables of the action of σ on indices.

³We use the word composition in a broader sense, given that, strictly speaking, a composition should not contain any zeroes. Similarly, we allow the presence of zeroes in a partition.

Finally, to any superpartition Λ , we associate a unique standard partition Λ^* obtained by rearranging the parts of the superpartition in decreasing order:

$$\Lambda^* = (\Lambda_c)^+ . \quad (14)$$

For instance, the $*$ -rearrangement of $(4, 2, 1; 5, 3, 2, 1)$ is

$$(4, 2, 1; 5, 3, 2, 1)^* = (5, 4, 3, 2, 2, 1, 1) . \quad (15)$$

A natural basis of P^{S_N} is provided by the monomial symmetric superfunctions⁴

$$m_\Lambda = \frac{1}{f_\Lambda} \sum_{\sigma \in S_N} \mathcal{K}_\sigma \theta_{\{1, \dots, m\}} z^\Lambda , \quad (16)$$

where the normalization constant f_Λ is

$$f_\Lambda = f_{\Lambda^s} = n_{\Lambda^s}(0)! n_{\Lambda^s}(1)! n_{\Lambda^s}(2)! \cdots , \quad (17)$$

with $n_{\Lambda^s}(i)$ the number of i 's in Λ^s , the symmetric part of the superpartition $\Lambda = (\Lambda^a; \Lambda^s)$. This normalization ensures that the coefficient of the monomial $\theta_{\{1, \dots, m\}} z^\Lambda$ appearing in the expansion of m_Λ is equal to 1. The supermonomial $m_{(3,1;2,1)}$ is given in (4) for $N = 4$.

Finally, we will define a scalar product $\langle \cdot, \cdot \rangle_\beta$ in P . With

$$\Delta(z) = \prod_{1 \leq j < k \leq N} \left[\frac{z_j - z_k}{z_j z_k} \right] , \quad (18)$$

$\langle \cdot, \cdot \rangle_\beta$ is defined (for β a positive integer) on the basis elements of P as

$$\langle \theta_I z^\lambda, \theta_J z^\mu \rangle_\beta = \begin{cases} \text{C.T.} [\Delta^\beta(z) \Delta^\beta(\bar{z}) z^\lambda / z^\mu] & \text{if } I = J \\ 0 & \text{otherwise} \end{cases} , \quad (19)$$

where $\bar{z}_i = 1/z_i$, and where C.T. $[E]$ stands for the constant term of the expression E . This scalar product is a special case of the physical scalar product of the underlying supersymmetric quantum many-body problem (β is now arbitrary)

$$\langle A(z, \theta), B(z, \theta) \rangle_\beta = \left(\prod_{i=1}^N \oint \frac{dz_i}{2\pi i z_i} \right) \int d\theta_1 \cdots d\theta_N \Delta^\beta(z) \Delta^\beta(\bar{z}) A(z, \theta), B(\bar{z}, \bar{\theta}) , \quad (20)$$

where $\overline{\theta_{i_1} \cdots \theta_{i_m}}$ is defined such that

$$(\theta_{i_1} \cdots \theta_{i_m}) (\overline{\theta_{i_1} \cdots \theta_{i_m}}) = \theta_N \cdots \theta_1 , \quad (21)$$

an operation akin to the Hodge duality transformation. For instance, if $N = 5$, we have $\overline{\theta_2 \theta_5} = -\theta_4 \theta_3 \theta_1$. The integral over fermionic variables refers to the Berezin integration

$$\int d\theta = 0 , \quad \int d\theta \theta = 1 . \quad (22)$$

⁴The quantity z^Λ is to be understood as $z_1^{\Lambda_1} \cdots z_N^{\Lambda_N}$, that is, as if Λ were replaced by Λ_c . However, to alleviate the notation, we will omit the subindex c when Λ is treated as a formal power.

3 Dunkl operators and conserved charges

The Dunkl operators, $\mathcal{D}_i$, are defined as [10]⁵

$$\begin{aligned}\mathcal{D}_i &= z_i \frac{\partial}{\partial z_i} + \beta \sum_{j < i} \frac{z_i}{z_i - z_j} (1 - K_{ij}) + \beta \sum_{j > i} \frac{z_j}{z_i - z_j} (1 - K_{ij}) - \beta(i - 1) \\ &= z_i \frac{\partial}{\partial z_i} + \beta \sum_{j < i} \mathcal{O}_{ij} + \beta \sum_{j > i} \mathcal{O}_{ij} - \beta(i - 1),\end{aligned}\tag{23}$$

where

$$\mathcal{O}_{ij} = \begin{cases} \frac{z_i}{z_i - z_j} (1 - K_{ij}) & j < i \\ \frac{z_j}{z_i - z_j} (1 - K_{ij}) & j > i \end{cases}.\tag{24}$$

The set $\{\mathcal{D}_i\}$ forms a family of commuting operators satisfying the degenerate Hecke relations

$$K_{i,i+1} \mathcal{D}_{i+1} - \mathcal{D}_i K_{i,i+1} = \beta \quad \text{and} \quad K_{j,j+1} \mathcal{D}_i = \mathcal{D}_i K_{j,j+1} \quad (i \neq j, j+1).\tag{25}$$

It turns out that any conserved charge, C_n , of the stCMS model can be written as the P^{S_N} -projection of an expression, $\mathcal{C}_n$, involving Dunkl operators

$$\mathcal{C}_n|_{P^{S_N}} = C_n \quad \Longleftrightarrow \quad \mathcal{C}_n f = C_n f, \quad \forall f \in P^{S_N}.\tag{26}$$

This is a key tool in our subsequent analysis, since by working with Dunkl operators, we avoid manipulating fermionic variables to a large extent. More explicitly, the stCMS conserved charges are defined as the projection onto P^{S_N} of the following expressions

$$\mathcal{H}_n = \sum_{i=1}^N \mathcal{D}_i^n,\tag{27}$$

$$\mathcal{Q}_n = \sum_{w \in S_N} \mathcal{K}_w (\theta_1 \mathcal{D}_1^n),\tag{28}$$

$$\mathcal{Q}_n^\dagger = \sum_{w \in S_N} \mathcal{K}_w \left(\frac{\partial}{\partial \theta_1} \mathcal{D}_1^n \right),\tag{29}$$

$$\mathcal{I}_n = \sum_{w \in S_N} \mathcal{K}_w \left(\theta_1 \frac{\partial}{\partial \theta_1} \mathcal{D}_1^n \right).\tag{30}$$

Of these charges, $\mathcal{I}_0/(N-1)!$ gives the fermion number. Observe that $\mathcal{H}_n$ and $\mathcal{I}_n$ preserve the number of fermions (the number of θ_i 's) of the superpolynomials on which they act, while $\mathcal{Q}_n$ (resp. $\mathcal{Q}_n^\dagger$) increases it by 1 (resp. -1). Also, since $\mathcal{H}_n$ is known to be central in the degenerate Hecke algebra [6], it commutes with K_σ , for any element σ of the symmetric group. Finally, these expressions not being unique, we should mention that the present choice for $\mathcal{H}_n$ and $\mathcal{I}_n$ is motivated by the requirement that they act triangularly on monomial superfunctions (as we will show later on).

We finish this section with two propositions, the first one relying on the following lemma.

Lemma 3. *For any nonnegative integers n and m , we have*

$$[\mathcal{D}_1^n \mathcal{K}_{12}, \mathcal{D}_1^m \mathcal{K}_{12}] + [\mathcal{K}_{12} \mathcal{D}_1^n, \mathcal{K}_{12} \mathcal{D}_1^m] = 0.\tag{31}$$

⁵Following [4], we use the qualitative ‘Dunkl’ for all Dunkl-type operators. The present $\mathcal{D}_i$ ’s are often called Cherednik operators.

Proof. Given that κ_{12} (in $\mathcal{K}_{12}$) commutes with $\mathcal{D}_1$, the lemma is equivalent to

$$[\mathcal{D}_1^n K_{12}, \mathcal{D}_1^m K_{12}] + [K_{12} \mathcal{D}_1^n, K_{12} \mathcal{D}_1^m] = 0. \quad (32)$$

We will now seek to prove this expression. First, it is easy to verify, using $K_{12} \mathcal{D}_1 = \mathcal{D}_2 K_{12} - \beta$, that

$$K_{12} \mathcal{D}_1^n = \mathcal{D}_2^n K_{12} - \beta(\mathcal{D}_1^{n-1} + \mathcal{D}_1^{n-2} \mathcal{D}_2 + \cdots \mathcal{D}_1 \mathcal{D}_2^{n-2} + \mathcal{D}_2^{n-1}) = \mathcal{D}_2^n K_{12} - \beta h_{n-1}(\mathcal{D}_1, \mathcal{D}_2), \quad (33)$$

where $h_i(x_1, x_2)$ is the homogenous symmetric function of degree i in the variables x_1 and x_2 .

Using (33) and the commutativity of $\mathcal{D}_1$ and $\mathcal{D}_2$, we obtain

$$\begin{aligned} [\mathcal{D}_1^n K_{12}, \mathcal{D}_1^m K_{12}] + [K_{12} \mathcal{D}_1^n, K_{12} \mathcal{D}_1^m] &= \mathcal{D}_1^n \mathcal{D}_2^m - \beta h_{m-1} \mathcal{D}_1^n K_{12} - \mathcal{D}_1^m \mathcal{D}_2^n + \beta h_{n-1} \mathcal{D}_1^m K_{12} \\ &\quad + \mathcal{D}_2^n \mathcal{D}_1^m - \beta h_{n-1} K_{12} \mathcal{D}_1^m - \mathcal{D}_2^m \mathcal{D}_1^n + \beta h_{m-1} K_{12} \mathcal{D}_1^n \\ &= -\beta h_{m-1} \mathcal{D}_1^n K_{12} + \beta h_{n-1} \mathcal{D}_1^m K_{12} - \beta h_{n-1} \mathcal{D}_2^m K_{12} + \beta^2 h_{n-1} h_{m-1} \\ &\quad + \beta h_{m-1} \mathcal{D}_2^n K_{12} - \beta^2 h_{n-1} h_{m-1} \\ &= \beta h_{m-1} (\mathcal{D}_2^n - \mathcal{D}_1^n) K_{12} + \beta h_{n-1} (\mathcal{D}_1^m - \mathcal{D}_2^m) K_{12}, \end{aligned} \quad (34)$$

where h_i stands for $h_i(\mathcal{D}_1, \mathcal{D}_2)$. Finally, using the simple identity $(x_1^m - x_2^m) = (x_1 - x_2) h_{m-1}(x_1, x_2)$, the previous expression vanishes, and (32) is thus seen to hold. $\square$

Proposition 4. *The families of operators $\mathcal{H}_n$, $n = 1, \dots, N$, and $\mathcal{I}_m$, $m = 1, \dots, N$, when acting on P^{S_N} , form a set of mutually commuting operators, that is, they satisfy*

$$[\mathcal{H}_n, \mathcal{H}_m]f = [\mathcal{H}_n, \mathcal{I}_m]f = [\mathcal{I}_n, \mathcal{I}_m]f = 0, \quad (35)$$

for any $f \in P^{S_N}$.

Proof. Since the Dunkl operators $\mathcal{D}_i$ mutually commute, we have immediately $[\mathcal{H}_n, \mathcal{H}_m] = 0$. Further, since $\mathcal{H}_n$ commutes with $\mathcal{K}_\sigma$ for any permutation σ , we also get $[\mathcal{H}_n, \mathcal{I}_m] = 0$. The relation $[\mathcal{I}_n, \mathcal{I}_m]f = 0$ is less trivial. We have

$$\begin{aligned} \mathcal{I}_n \mathcal{I}_m f &= \sum_{w, \sigma \in S_N} \mathcal{K}_w \theta_1 \frac{\partial}{\partial \theta_1} \mathcal{D}_1^n \mathcal{K}_\sigma \theta_1 \frac{\partial}{\partial \theta_1} \mathcal{D}_1^m f \\ &= \sum_{w, \sigma \in S_N} \theta_{(w)_1} \frac{\partial}{\partial \theta_{(w)_1}} \theta_{(w\sigma)_1} \frac{\partial}{\partial \theta_{(w\sigma)_1}} \mathcal{K}_w \mathcal{D}_1^n \mathcal{K}_\sigma \mathcal{D}_1^m f, \end{aligned} \quad (36)$$

where $(w)_1$ is the first entry of the permutation w . Therefore, $[\mathcal{I}_n, \mathcal{I}_m]f$ can be written as

$$[\mathcal{I}_n, \mathcal{I}_m]f = \sum_{w, \sigma \in S_N} \theta_{(w)_1} \frac{\partial}{\partial \theta_{(w)_1}} \theta_{(w\sigma)_1} \frac{\partial}{\partial \theta_{(w\sigma)_1}} (\mathcal{K}_w \mathcal{D}_1^n \mathcal{K}_\sigma \mathcal{D}_1^m - \mathcal{K}_w \mathcal{D}_1^m \mathcal{K}_\sigma \mathcal{D}_1^n) f. \quad (37)$$

To prove that this expression is equal to zero, we will match its summands (w, σ) and $(w\sigma, \sigma^{-1})$, and see that they cancel each others. First, if we let $w \rightarrow w\sigma$, and $\sigma \rightarrow \sigma^{-1}$, the summand (w, σ) of (37) becomes

$$\begin{aligned} \theta_{(w\sigma)_1} \frac{\partial}{\partial \theta_{(w\sigma)_1}} \theta_{(w)_1} \frac{\partial}{\partial \theta_{(w)_1}} (\mathcal{K}_w \mathcal{K}_\sigma \mathcal{D}_1^n \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^m - \mathcal{K}_w \mathcal{K}_\sigma \mathcal{D}_1^m \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^n) f \\ = \theta_{(w)_1} \frac{\partial}{\partial \theta_{(w)_1}} \theta_{(w\sigma)_1} \frac{\partial}{\partial \theta_{(w\sigma)_1}} (\mathcal{K}_w \mathcal{K}_\sigma \mathcal{D}_1^n \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^m - \mathcal{K}_w \mathcal{K}_\sigma \mathcal{D}_1^m \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^n) f, \end{aligned} \quad (38)$$

the equality being obtained by interchanging the two prefactors $\theta_i \partial / \partial \theta_i$. Now, $(w, \sigma) = (w\sigma, \sigma^{-1})$ iff $\sigma = e$. Since, in the case $(w, \sigma) = (w, e)$, the summand (w, σ) of (37) cancels, we can assume that (w, σ) and

$(w\sigma, \sigma^{-1})$ are distinct summands. Having shown that the prefactors are the same for the two summands (w, σ) and $(w\sigma, \sigma^{-1})$ (cf. (38)), verifying their cancellation amounts to checking that

$$\left(\mathcal{K}_w \mathcal{D}_1^n \mathcal{K}_\sigma \mathcal{D}_1^m - \mathcal{K}_w \mathcal{D}_1^m \mathcal{K}_\sigma \mathcal{D}_1^n \right) f + \left(\mathcal{K}_w \mathcal{K}_\sigma \mathcal{D}_1^n \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^m - \mathcal{K}_w \mathcal{K}_\sigma \mathcal{D}_1^m \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^n \right) f = 0. \quad (39)$$

Since $f \in P^{S_N}$, this is equivalent to

$$\left(\mathcal{D}_1^n \mathcal{K}_\sigma \mathcal{D}_1^m \mathcal{K}_{\sigma^{-1}} - \mathcal{D}_1^m \mathcal{K}_\sigma \mathcal{D}_1^n \mathcal{K}_{\sigma^{-1}} + \mathcal{K}_\sigma \mathcal{D}_1^n \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^m - \mathcal{K}_\sigma \mathcal{D}_1^m \mathcal{K}_{\sigma^{-1}} \mathcal{D}_1^n \right) f = 0. \quad (40)$$

Now, $\mathcal{D}_1$ commutes with $\mathcal{K}_{i, i+1}$, as long as $i \neq 1$. Therefore, if σ leaves 1 fixed, (40) holds. We can thus assume that σ does not leave 1 fixed. In this case, σ can be decomposed as $\alpha(12)\beta$, where α and β are permutations that leave 1 invariant, and (40) becomes

$$\mathcal{K}_\alpha \left(\mathcal{D}_1^n \mathcal{K}_{12} \mathcal{D}_1^m \mathcal{K}_{12} - \mathcal{D}_1^m \mathcal{K}_{12} \mathcal{D}_1^n \mathcal{K}_{12} + \mathcal{K}_{12} \mathcal{D}_1^n \mathcal{K}_{12} \mathcal{D}_1^m - \mathcal{K}_{12} \mathcal{D}_1^m \mathcal{K}_{12} \mathcal{D}_1^n \right) \mathcal{K}_{\alpha^{-1}} f = 0, \quad (41)$$

where we have used the facts that $\mathcal{K}_{12}^{-1} = \mathcal{K}_{12}$, and that $\mathcal{K}_\alpha$ and $\mathcal{K}_\beta$ commute with $\mathcal{D}_1$. Given that from Lemma 3, $[\mathcal{D}_1^n \mathcal{K}_{12}, \mathcal{D}_1^m \mathcal{K}_{12}] + [\mathcal{K}_{12} \mathcal{D}_1^n, \mathcal{K}_{12} \mathcal{D}_1^m] = 0$, the expression is finally seen to hold, thereby proving $[\mathcal{I}_n, \mathcal{I}_m]f = 0$. $\square$

Proposition 5. *The charges $\mathcal{H}_n$ and $\mathcal{I}_n$ are self-adjoint with respect to the scalar product (19).*

Proof. In the $\mathcal{H}_n$ case, this simply follows from the self-adjointness of the operators $\mathcal{D}_i$ [6]. In the case of $\mathcal{I}_n$, we also need to use $\theta_1^\dagger = \frac{\partial}{\partial \theta_1}$, and

$$\left(\theta_1 \frac{\partial}{\partial \theta_1} \right)^\dagger = \left(\frac{\partial}{\partial \theta_1} \right)^\dagger \theta_1^\dagger = \theta_1 \frac{\partial}{\partial \theta_1}. \quad (42)$$

$\square$

Our first goal will be to find the common eigenfunctions of the commuting operators $\mathcal{H}_n$ and $\mathcal{I}_n$. However, before plunging into the relevant computations, we need to introduce further technical tools. This will be the subject of the following two sections.

4 Orderings on superpartitions

In this section we introduce three orderings on superpartitions. They will provide three different ways of defining triangular decompositions.

First recall the usual dominance ordering on partitions [3]. If λ and μ are two partitions (i.e., $\lambda = \lambda^+$ and $\mu = \mu^+$), then $\lambda \geq \mu$ iff $\lambda_1 + \dots + \lambda_i \geq \mu_1 + \dots + \mu_i$ for all i . This ordering can be extended to compositions as follows. Any composition λ can be obtained from its rearranged partition λ^+ by a sequence of permutations. Among all permutations w such that $\lambda = w\lambda^+$, there exists a unique one, denoted w_λ , of minimal length.

Definition 6. Given two compositions λ, μ , we say that $\lambda \geq \mu$ if either $\lambda^+ > \mu^+$, or $\lambda^+ = \mu^+$ and $w_\lambda \leq w_\mu$ in the Bruhat order of the symmetric group. This will be called the *Bruhat ordering on compositions*.⁶

An immediate consequence of this definition is that $\lambda^+ \geq \mu$ for any composition μ such that $\mu^+ = \lambda^+$. Moreover, given that to any superpartition is associated a composition, this ordering induces an ordering on superpartitions.

⁶The ordering on compositions could alternatively be formulated as follows [8]. We say that $\lambda \geq \mu$ if either $\lambda^+ > \mu^+$, or $\lambda^+ = \mu^+$ and $\sum_{i=1}^k \lambda_i \geq \sum_{i=1}^k \mu_i$ for all k .

Definition 7. Given two superpartitions Λ, Ω , we say that $\Lambda \geq \Omega$, if $\Lambda_c \geq \Omega_c$. We shall refer to this ordering as the *Bruhat ordering on superpartitions*.

We finally define two other orderings on superpartitions.

Definition 8. The *h-ordering* $\leq_h$ is defined such that $\Omega \leq_h \Lambda$, if either $\Lambda = \Omega$, or $\Omega^* \neq \Lambda^*$ and $\Omega \leq \Lambda$.

Definition 9. The *t-ordering* $\leq_t$ is defined such that $\Omega \leq_t \Lambda$, iff $\Lambda^* = \Omega^*$ and $\Omega \leq \Lambda$.

Obviously, these two new orderings on superpartitions are special cases of the Bruhat ordering in the sense that if either $\Omega \leq_h \Lambda$ or $\Omega \leq_t \Lambda$, then $\Omega \leq \Lambda$.

Let us look at illustrative examples. The two superpartitions $\Omega = (5, 3; 4, 1, 1)$ and $\Lambda = (5, 1; 4, 4, 0)$ cannot be *t*-compared since $\Omega^* \neq \Lambda^*$. However, they can be *h*-compared since $(\Omega_c)^+ = \Omega^* = (5, 4, 3, 1, 1) < (\Lambda_c)^+ = \Lambda^* = (5, 4, 4, 1, 0)$. On the other hand, let us see how $\Gamma = (5, 1; 4, 3, 1)$ compares with the previous two superpartitions. Again, Γ and Λ cannot be *t*-compared, but are such that $\Gamma <_h \Lambda$. Since $\Gamma^* = \Omega^*$, the two superpartitions Γ and Ω may be *t*-comparable. With $\Gamma_c = (5, 1, 4, 3, 1) = \sigma_2 \sigma_3(5, 4, 3, 1, 1)$ and $\Omega_c = (5, 3, 4, 1, 1) = \sigma_2(5, 4, 3, 1, 1)$, we in fact conclude that $\Gamma <_t \Omega$.

As we will see later, the *h*-ordering characterizes the triangular action of the charges $\mathcal{H}_n$ on the supermonomial basis, thus justifying its labeling *h*.⁷ On the other hand, the action of the charges $\mathcal{I}_n$ on the non-orthogonal Jack superpolynomials will be shown to be triangular with respect to a different ordering. This will happen to be the *t*-ordering. The label *t* refers in this case to the interchange operator *T*, which we introduce below, relating the superpartitions that can be *t*-compared.

The rest of this section is devoted to presenting some properties of the Bruhat ordering on compositions. Remarkably, the action of a Dunkl operator $\mathcal{D}_i$ is triangular with respect to this ordering.

Property 10. [6, 7]. Let λ be a composition. Then,

$$\mathcal{D}_i z^\lambda = \underline{\lambda}_i z^\lambda + \sum_{\mu < \lambda} c_{\lambda\mu} z^\mu, \quad (43)$$

where

$$\underline{\lambda}_i = \lambda_i - \beta \left(\#\{j = 1, \dots, i-1 \mid \lambda_j \geq \lambda_i\} + \#\{j = i+1, \dots, N \mid \lambda_j > \lambda_i\} \right). \quad (44)$$

For $i < j$, let T_{ij} be such that on a composition λ ,

$$T_{ij}\lambda = \begin{cases} (\dots \lambda_j \dots \lambda_i \dots) & \text{if } \lambda_i > \lambda_j \\ (\dots \lambda_i \dots \lambda_j \dots) & \text{otherwise} \end{cases}, \quad (45)$$

that is, T_{ij} interchanges the entries λ_i and λ_j only when $\lambda_i > \lambda_j$. The action of T_{ij} on superpartitions Λ is defined via the corresponding compositions Λ_c . The order on compositions satisfies the following obvious property.

Property 11. Let μ and λ be two compositions such that $\mu \leq \lambda$, and $\mu^+ = \lambda^+$, that is, such that $\mu \leq_t \lambda$. Then, there exists a sequence of operators T_{ij} giving

$$\mu = T_{i_1 j_1} \cdots T_{i_\ell j_\ell} \lambda. \quad (46)$$

Lemma 12. Let Ω and Λ be two superpartitions. If

$$\Omega = \overline{T_{i_1 j_1} \cdots T_{i_\ell j_\ell} \Lambda}, \quad (47)$$

for some $T_{i_k j_k}$, $1 \leq k \leq \ell$, then $\Omega \leq \Lambda$.

⁷This is the ordering introduced in [4]. In reference [5], a more precise formulation of this ordering was introduced (and called $\leq_s$).

First, it is important to realize that the product $T_{i_1 j_1} \cdots T_{i_\ell j_\ell}$ can be rewritten as a product of T_{ij} 's where all the T_{ij} 's that interchange elements between the fermionic and bosonic sectors (that is, that interchange entries of Λ^a and Λ^s) are to the right. Since, in this form, the remaining elements only interchange entries within each sectors, their action will amount to nothing after the 'bar' operation has been performed. We can therefore assume that all the T_{ij} 's in $T_{i_1 j_1} \cdots T_{i_\ell j_\ell}$ interchange elements between the fermionic and bosonic sectors.

Before going into the proof of the lemma, let us first give an example that will hopefully shed some light on the many steps involved in the proof. Let us consider the superpartition

$$\Lambda = (7, 5, 4, 3, 0; 9, 6, 4, 4, 2, 2, 1, 1, 1), \quad (48)$$

and act on it with $T_{1,11}$, and then with $T_{4,13}$. We have thus

$$T_{4,13}T_{1,11}\Lambda = T_{4,13}T_{1,11}(7, 5, 4, 3, 0; 9, 6, 4, 4, 2, 2, 1, 1, 1) = (2, 5, 4, 1, 0; 9, 6, 4, 4, 2, 7, 1, 3, 1). \quad (49)$$

The superpartition Ω is obtained by applying the 'bar' operation:

$$\Omega = \overline{(2, 5, 4, 1, 0; 9, 6, 4, 4, 2, 7, 1, 3, 1)} = (5, 4, 2, 1, 0; 9, 7, 6, 4, 4, 3, 2, 1, 1). \quad (50)$$

Now, can we conclude directly that $\Omega < \Lambda$? No, because even though Λ_c Bruhat dominates the intermediate composition $(2, 5, 4, 1, 0; 9, 6, 4, 4, 2, 7, 1, 3, 1)$, the latter does not dominate (it is actually dominated by) the composition Ω_c associated to the resulting superpartition Ω . In the intermediate step, we somehow ended up too low to apply a chain of Bruhat dominance. This simply indicates that the 'bar' operation is not compatible with the ordering on compositions. Therefore, the lemma does not follow immediately from the previous property. Actually, what the proof of the lemma gives is a precise construction to arrive at Ω via a sequence of T_{ij} 's without introducing rearrangements at any intermediate step.

Proof: Essentially, we want to show that any Ω that can be obtained from Λ by exchanging a certain number of elements of Λ^a and Λ^s , and then rearranging both vectors, can also be obtained by simply applying a sequence of T_{ij} 's, without rearrangement. Let $(a_1, \dots, a_\ell)$ be the partition corresponding to the elements of Λ^a that will be moved to the symmetric side. Also, let $(p_1, \dots, p_\ell)$ be their respective positions in Λ , and $(p'_1, \dots, p'_\ell)$ be their final positions, that is, their positions in Ω . Similarly, let $(b_1, \dots, b_\ell)$ be the partition corresponding to the elements of Λ^s that will be moved to the antisymmetric side, and denote by $(q_1, \dots, q_\ell)$ their positions in Λ , and by $(q'_1, \dots, q'_\ell)$ their final positions in Ω . Because we move larger elements to the symmetric side, we must have $a_k > b_k$ for all $k = 1, \dots, \ell$. In our example, we have $\ell = 2$, and

$$(a_1, a_2) = (7, 3) \quad (p_1, p_2) = (1, 4) \quad (p'_1, p'_2) = (7, 11) \quad (51)$$

$$(b_1, b_2) = (2, 1) \quad (q_1, q_2) = (10, 12) \quad (q'_1, q'_2) = (3, 4). \quad (52)$$

Now, start from Λ and move $(a_1, \dots, a_\ell)$ so that they occupy the intermediate positions $q'_1, \dots, q'_\ell$ respectively. This can be done using a sequence of T_{ij} because, from $a_k > b_k$, we know that all the a_k 's are moved to the right passed smaller elements. The precise sequence of T_{ij} 's that performs this operation is $T_{p_1 q'_1} \cdots T_{p_\ell q'_\ell}$. In the resulting vector, move $(b_1, \dots, b_\ell)$ so that they occupy positions $q_1, \dots, q_\ell$ respectively. Again this can be done using T_{ij} operators because, from $a_k > b_k$ and choosing b_m such that it occupies the leftmost position whenever there are multiplicities, all the a_k 's will be moved to the left passed larger elements. This amounts to applying $T_{p'_1 q_1} \cdots T_{p'_\ell q_\ell}$. Finally, applying the sequence $T_{p'_1 q'_1} \cdots T_{p'_\ell q'_\ell}$ gives Ω . Transposing these various steps to our example yields

$$T_{3,7}T_{4,11}T_{7,10}T_{11,12}T_{1,3}T_{4,4}(7, 5, 4, 3, 0; 9, 6, 4, 4, 2, 2, 1, 1, 1) = (5, 4, 2, 1, 0; 9, 7, 6, 4, 4, 3, 2, 1, 1) = \Omega \quad (53)$$

This shows that $\Omega \leq \Lambda$. □

Corollary 13. Let $\mu = (\mu_1, \dots, \mu_m; \mu_{m+1}, \dots, \mu_N)$ be such that $\bar{\mu} = \Omega$. If $\Omega \leq \Lambda$, then $\overline{T_{ij} \mu} \leq \Lambda$.

Proof: Since $\bar{\mu} = \Omega$, μ can be written as $\mu = T_{i_1 j_1} \cdots T_{i_\ell j_\ell} \Omega$, for some operators $T_{i_k j_k}$, $k = 1, \dots, \ell$. Therefore, from Lemma 12,

$$\overline{T_{ij} \mu} = \overline{T_{ij} T_{i_1 j_1} \cdots T_{i_\ell j_\ell} \Omega} \leq \Omega, \quad (54)$$

which gives $\overline{T_{ij} \mu} \leq \Lambda$ if $\Omega \leq \Lambda$. $\square$

5 Triangular Operators and Determinants

This section presents basic results regarding triangular operators. We should point out that Theorem 16 and Corollary 17 appear for instance in a disguised form in [3]. The exposition of the material in this section follows that of [11, 12].

Let $\{s_\Lambda\}_\Lambda$ be any basis of P^{S_N} . We write $P_{\Lambda, \preceq}^{(s)}$ for the finite-dimensional subspace of P^{S_N} spanned by the s_Ω 's such that $\Omega \preceq \Lambda$, with respect to some ordering $\preceq$ (which could be any of the three orderings introduced previously), i.e.,

$$P_{\Lambda, \preceq}^{(s)} = \text{Span}\{s_\Omega\}_{\Omega \preceq \Lambda}. \quad (55)$$

Definition 14. A linear operator $O_t : P^{S_N} \rightarrow P^{S_N}$ is called *triangular* if $O_t(P_{\Lambda, \preceq}^{(s)}) \subseteq P_{\Lambda, \preceq}^{(s)}$ for every superpartition Λ .

The triangularity of a linear operator O_t in P^{S_N} reduces its eigenvalue problem to a finite-dimensional one. Triangular operators can be diagonalized through a determinantal representation of the eigenfunctions. The triangularity implies that the expansion of $O_t s_\Lambda$ is of the form

$$O_t s_\Lambda = \epsilon_\Lambda s_\Lambda + \sum_{\Omega \prec \Lambda} d_{\Lambda\Omega} s_\Omega, \quad (56)$$

with the diagonal matrix elements ϵ_Λ being precisely the eigenvalues of O_t .

Definition 15. The triangular operator O_t is called *regular* if $\epsilon_\Omega \neq \epsilon_\Lambda$ whenever $\Omega \prec \Lambda$.

Let $\{p_\Lambda\}_\Lambda$ be a corresponding basis of eigenfunctions diagonalizing O_t . Clearly, we can choose p_Λ to have an expansion of the form

$$p_\Lambda = s_\Lambda + \sum_{\Omega \prec \Lambda} c_{\Lambda\Omega} s_\Omega, \quad (57)$$

where the normalization has been chosen to make p_Λ monic. The following theorem provides an explicit determinantal formula for p_Λ , given the action of O_t on s_Λ expressed in the basis s_Λ .

Theorem 16. Let O_t be a regular triangular operator in P^{S_N} whose action on the basis $\{s_\Lambda\}_\Lambda$ is given by (56). Then the unique monic basis $\{p_\Lambda\}_\Lambda$ of P^{S_N} triangularly related to the basis $\{s_\Lambda\}_\Lambda$ (cf. (57)) diagonalizing O_t , i.e.,

$$O_t p_\Lambda = \epsilon_\Lambda p_\Lambda, \quad \forall \Lambda, \quad (58)$$

is given explicitly by the (lower) Hessenberg determinant

$$p_\Lambda = \frac{1}{\mathcal{E}_\Lambda} \begin{vmatrix} s_{\Lambda(1)} & \epsilon_{\Lambda(1)} - \epsilon_{\Lambda(n)} & 0 & \cdots & \cdots & 0 \\ s_{\Lambda(2)} & d_{\Lambda(2)\Lambda(1)} & \epsilon_{\Lambda(2)} - \epsilon_{\Lambda(n)} & 0 & \cdots & 0 \\ \vdots & \vdots & & \ddots & \ddots & \vdots \\ \vdots & \vdots & \vdots & & \ddots & 0 \\ s_{\Lambda(n-1)} & d_{\Lambda(n-1)\Lambda(1)} & d_{\Lambda(n-1)\Lambda(2)} & \cdots & & \epsilon_{\Lambda(n-1)} - \epsilon_{\Lambda(n)} \\ s_{\Lambda(n)} & d_{\Lambda(n)\Lambda(1)} & d_{\Lambda(n)\Lambda(2)} & \cdots & \cdots & d_{\Lambda(n)\Lambda(n-1)} \end{vmatrix}. \quad (59)$$

Here $\Lambda^{(1)} < \Lambda^{(2)} < \dots < \Lambda^{(n-1)} < \Lambda^{(n)} = \Lambda$ denotes any linear ordering, refining the natural order $\preceq$, of the superpartitions, $\Lambda^{(i)}, i = 1, \dots, n-1$, that precede Λ in the ordering $\preceq$. The normalization is determined by

$$\mathcal{E}_\Lambda = \prod_{i=1}^{n-1} (\epsilon_\Lambda - \epsilon_{\Lambda^{(i)}}). \quad (60)$$

With $O_t = \mathcal{H}_2$ and $s_\Lambda = m_\Lambda$, the previous theorem leads to a closed expression for the $\mathcal{H}_2$ eigenfunctions, the Jack superpolynomials $\mathcal{J}_\Lambda$ of [4, 5], in terms of the coefficients $d_{\Lambda\Omega}$ entering in the supermonomial decomposition of $\mathcal{H}_2 m_\Lambda$. These coefficients have been computed in [5]. As already indicated, the superpolynomials $\mathcal{J}_\Lambda$ are not orthogonal. We will seek linear combinations that are orthogonal by considering the eigenfunctions of $\mathcal{I}_1$. Theorem 16 will then be invoked again, but this time with $s_\Lambda = \mathcal{J}_\Lambda$ and $O_t = \mathcal{I}_1$. Computing the action of $\mathcal{I}_1$ in the $\mathcal{J}_\Lambda$ basis will provide closed form formulas for the orthogonal superpolynomials J_Λ . In the m_Λ basis, J_Λ will appear as a determinant of determinants.

As an aside, we point out that the determinantal formula for p_Λ , leads to a linear recurrence relation encoding an efficient algorithm for the computation of the coefficients $c_{\Lambda\Omega}$ entering the expansion (57).

Corollary 17. *The expansion of p_Λ is of the form*

$$p_\Lambda = \sum_{\ell=1}^n c_{\Lambda\Lambda^{(\ell)}} s_{\Lambda^{(\ell)}}, \quad (61)$$

with $c_{\Lambda\Lambda^{(n)}} = c_{\Lambda\Lambda} = 1$ and, for $1 < \ell \leq n$,

$$c_{\Lambda\Lambda^{(\ell-1)}} = \frac{1}{\epsilon_\Lambda - \epsilon_{\Lambda^{(\ell-1)}}} \sum_{k=\ell}^n c_{\Lambda\Lambda^{(k)}} d_{\Lambda^{(k)}\Lambda^{(\ell-1)}}. \quad (62)$$

We conclude this section with an elementary and surely well known proposition that we prove for a lack of reference. It provides a simple way of computing the p_Λ eigenvalues of mutually commuting operators in terms of the action of these operators on the s_Λ basis.

Proposition 18. *Let D_t be a triangular operator commuting with O_t . Then,*

$$D_t p_\Lambda = \varepsilon_\Lambda p_\Lambda, \quad (63)$$

where ε_Λ is the coefficient of s_Λ in $D_t s_\Lambda$.

Proof. Let $\tilde{p}_\Lambda = D_t p_\Lambda / \varepsilon_\Lambda$. Then, from (57) and the fact that D_t is a triangular operator, $\tilde{p}_\Lambda$ is seen to be of the form

$$\tilde{p}_\Lambda = s_\Lambda + \sum_{\Omega \preceq \Lambda} g_{\Lambda\Omega} s_\Omega. \quad (64)$$

Now, because O_t and D_t commute, we have $O_t \tilde{p}_\Lambda = D_t O_t p_\Lambda / \varepsilon_\Lambda = \epsilon_\Lambda \tilde{p}_\Lambda$. Therefore, the monic polynomial $\tilde{p}_\Lambda$ diagonalizes O_t and, from (64), is triangularly related to the basis $\{s_\Lambda\}_\Lambda$. From the uniqueness in Theorem 16, we must have $\tilde{p}_\Lambda = p_\Lambda$, or $D_t p_\Lambda = \varepsilon_\Lambda p_\Lambda$. $\square$

6 The action of $\mathcal{H}_n$

We are now ready to tackle one of our main objectives, which is to obtain common eigenfunctions of the commuting operators $\mathcal{H}_n$ and $\mathcal{I}_n$. In this section, we first study the action of the $\mathcal{H}_n$'s.

We start with a very simple proposition concerning the operators $\mathcal{O}_{ij}$ that we state without proof.

Proposition 19. *If we only consider terms that are permutations of $z_i^{\lambda_i} z_j^{\lambda_j}$, we have, for $i > j$,*

$$\mathcal{O}_{ij} z_i^{\lambda_i} z_j^{\lambda_j} = \begin{cases} z_i^{\lambda_i} z_j^{\lambda_j} & \lambda_i > \lambda_j \\ -z_i^{\lambda_j} z_j^{\lambda_i} & \lambda_i < \lambda_j \\ 0 & \text{otherwise} \end{cases}, \quad (65)$$

and, for $i < j$,

$$\mathcal{O}_{ij} z_i^{\lambda_i} z_j^{\lambda_j} = \begin{cases} z_i^{\lambda_j} z_j^{\lambda_i} & \lambda_i > \lambda_j \\ -z_i^{\lambda_i} z_j^{\lambda_j} & \lambda_i < \lambda_j \\ 0 & \text{otherwise} \end{cases}. \quad (66)$$

Lemma 20. *Let λ be a partition, and let λ^R be λ in reverse order. Then*

$$\mathcal{H}_n z^{\lambda^R} = \varepsilon_{n,\lambda} z^{\lambda^R} + \sum_{\mu < \lambda^R; \mu^+ \neq \lambda} a_{n,\mu} z^\mu, \quad (67)$$

with $\varepsilon_{n,\lambda}$ given explicitly by the formula

$$\varepsilon_{n,\lambda} = \sum_{i=1}^N \left(\underline{\lambda_i^R} \right)^n, \quad (68)$$

where the symbol $\underline{\gamma_i}$ was introduced in Property 10, and where λ_i^R stands for $(\lambda^R)_i$.

Proof. The lemma will hold if we can demonstrate that, for $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$, we have

$$\mathcal{D}_i z^{\lambda^R} = \underline{\lambda_i^R} z^{\lambda^R} + \sum_{\mu < \lambda^R; \mu^+ \neq \lambda} a_\mu z^\mu. \quad (69)$$

Using Property 10, for this to be true we only need to show that terms of the type z^μ , where $\mu^+ = \lambda$ never occur (except for z^{λ^R}). Since $\lambda^R \leq \mu$ for any μ such that $\mu^+ = \lambda$, this is indeed seen to be true. $\square$

The special action of $\mathcal{H}_n$ on z^{λ^R} induces a triangularity on m_Λ .

Theorem 21. *Let $\Lambda^* = (\lambda_1, \dots, \lambda_N) = \lambda$. Then,*

$$\mathcal{H}_n m_\Lambda = \varepsilon_{n,\lambda} m_\Lambda + \sum_{\Omega <_h \Lambda} a_{\Lambda\Omega}^{(n)} m_\Omega, \quad (70)$$

with $\varepsilon_{n,\lambda}$ given in Lemma 20.

Proof. We have that $\theta_{\{1,\dots,m\}} z^\Lambda = \pm \mathcal{K}_{\sigma'} \theta_I z^{\lambda^R}$, for some $\sigma' \in S_N$ and some $I \subseteq \{1, \dots, N\}$. Since $\mathcal{H}_n$ commutes with $\mathcal{K}_{ij}$ and θ_I , we therefore obtain, using equation (16) and Lemma 20,

$$\begin{aligned} \mathcal{H}_n m_\Lambda &= \pm \frac{1}{f_\Lambda} \sum_{\sigma \in S_N} \mathcal{K}_\sigma \mathcal{K}_{\sigma'} \theta_I \mathcal{H}_n z^{\lambda^R} \\ &= \pm \frac{1}{f_\Lambda} \sum_{\sigma \in S_N} \mathcal{K}_\sigma \mathcal{K}_{\sigma'} \theta_I \left(\varepsilon_{n,\lambda} z^{\lambda^R} + \sum_{\mu < \lambda^R; \mu^+ \neq \lambda} a_{n,\mu} z^\mu \right) \\ &= \varepsilon_{n,\lambda} m_\Lambda + \sum_{\Omega <_h \Lambda} a_{\Lambda\Omega}^{(n)} m_\Omega, \end{aligned}$$

which proves the theorem. $\square$

The next theorem was proven in [5] (cf. Theorem 10 therein).

Theorem 22. *There exists a unique basis $\{\mathcal{J}_\Lambda\}_\Lambda$ of P^{S_N} such that*

$$\mathcal{H}_2 \mathcal{J}_\Lambda = \varepsilon_{\lambda,2} \mathcal{J}_\Lambda \quad \text{and} \quad \mathcal{J}_\Lambda = m_\Lambda + \sum_{\Omega <_h \Lambda} v_{\Lambda\Omega} m_\Omega, \quad (71)$$

where $\lambda = \Lambda^*$.

The explicit action of $\mathcal{H}_2$ on the supermonomial basis, m_Λ , was computed in [5]. In view of Theorem 16, a determinantal formula giving $\mathcal{J}_\Lambda$ in terms of supermonomials immediately followed. It should be noted that an efficient algorithm to evaluate such a determinant can be found in Corollary 17.

We now show that $\mathcal{J}_\Lambda$ is an eigenfunction of $\mathcal{H}_n$, for all n .

Theorem 23. *With $\Lambda^* = \lambda$, we have*

$$\mathcal{H}_n \mathcal{J}_\Lambda = \varepsilon_{\lambda,n} \mathcal{J}_\Lambda = \left[\sum_{i=1}^N (\lambda_i^R)^n \right] \mathcal{J}_\Lambda. \quad (72)$$

Furthermore, for μ a partition such that $\mu \neq \lambda$, there exists at least one n such that $\varepsilon_{\lambda,n} \neq \varepsilon_{\mu,n}$.

Proof. Given Theorem 22, formula (70) and the fact that $\mathcal{H}_n$ and $\mathcal{H}_2$ commute, the first part of the theorem follows immediately from Proposition 18. The second part of the theorem is obvious because the $\varepsilon_{\lambda,n}$ ($n = 1, 2, \dots$) are polynomials in β whose constant terms $\varepsilon_{\lambda,n}|_{\beta=0} = \sum_{i=1}^N \lambda_i^n$ are such that if $\varepsilon_{\lambda,n}|_{\beta=0} = \varepsilon_{\mu,n}|_{\beta=0}$ for all $n = 1, 2, \dots$, then $\lambda = \mu$. Therefore, $\varepsilon_{\lambda,n}$ and $\varepsilon_{\mu,n}$, considered as functions of a generic parameter β , cannot be equal. $\square$

The last theorem implies that the superpolynomials $\mathcal{J}_\Lambda$ and $\mathcal{J}_\Omega$ associated to distinct superpartitions such that $\Lambda^* = \Omega^*$, share the same eigenvalues. Therefore, additional commuting operators need to be diagonalized in order to lift the degeneracies. These are the $\mathcal{I}_n$ charges, whose action is considered in the following section.

7 The action of $\mathcal{I}_n$

Let $\mathcal{I}'_n$ be the operator

$$\mathcal{I}'_n = \sum_{i=1}^N \mathcal{K}_{1i} \theta_1 \frac{\partial}{\partial \theta_1} \mathcal{D}_1^n. \quad (73)$$

The following proposition states that $\mathcal{I}_n$ and $\mathcal{I}'_n$ are equivalent (up to a constant) on P^{S_N} .

Proposition 24. *We have, for $f \in P^{S_N}$,*

$$\mathcal{I}_n f = (N-1)! \mathcal{I}'_n f. \quad (74)$$

Proof: The symmetric group can be factorized in the following way

$$\sum_{\sigma \in S_N} \mathcal{K}_\sigma = \sum_{i=1}^N \mathcal{K}_{1i} \sum_{w \in S_{\{2, \dots, N\}}} \mathcal{K}_w. \quad (75)$$

Therefore, since for $w \in S_{\{2, \dots, N\}}$, $\mathcal{K}_w$ commutes with $\theta_1 \frac{\partial}{\partial \theta_1} \mathcal{D}_1^n$, and since $\mathcal{K}_w$ leaves f invariant, the proposition follows. $\square$

We now introduce a subspace of P^{S_N} .

Definition 25. For Λ a superpartition in the m -fermion sector, L_Λ is given by

$$L_\Lambda = \text{Span} \{ z^\mu \mid \bar{\mu} \leq \Lambda \} = \text{Span} \{ K_\sigma K_w z^\Omega \mid \sigma \in S_m, w \in S_{\{m+1, \dots, N\}}, \text{ and } \Omega \leq \Lambda \},$$

where as usual Ω is a superpartition.

This subspace has the following property.

Lemma 26. *Let Λ be a superpartition in the m -fermion sector, and let $i \in \{1, \dots, m\}$. Then,*

$$\mathcal{D}_i(L_\Lambda) \subseteq L_\Lambda. \quad (76)$$

Proof. From the action of $\mathcal{D}_i$ given in Property 10, the lemma is false only if, for $\bar{\mu} \leq \Lambda$, there exist, in $\mathcal{D}_i z^\mu$, some terms of the type z^ν , where $\nu^+ = \Lambda^*$ and $\nu \not\leq \Lambda$. Given that by definition, $\mathcal{D}_i$ decomposes into the blocks $\mathcal{O}_{ij}$, it is thus sufficient to limit ourselves to the terms of $\mathcal{O}_{ij} z^\mu$ considered in Proposition 19. If $j \in \{1, \dots, m\}$, these terms are seen to be of the type z^ν , with $\bar{\nu} = \bar{\mu} \leq \Lambda$, and hence belong to L_Λ . On the other hand, when $j \in \{m+1, \dots, N\}$, since $i \in \{1, \dots, m\}$, we are always in the case $j > i$ of Proposition 19, in which case these terms are of the type $z^{T_{ij}\mu}$. The lemma then follows because, from Corollary 13, $\overline{T_{ij}\mu} \leq \Lambda$ whenever $\bar{\mu} \leq \Lambda$. $\square$

Theorem 27. *We have*

$$\mathcal{I}_n m_\Lambda = \epsilon_{\Lambda, n} m_\Lambda + \sum_{\Omega < \Lambda} b_{\Lambda\Omega}^{(n)} m_\Omega. \quad (77)$$

Proof: We will prove the equivalent statement that $\mathcal{I}'_n$ (see Proposition 24) acts triangularly. We have

$$\mathcal{I}'_n m_\Lambda = \sum_{i=1}^N \mathcal{K}_{1i} \theta_1 \frac{\partial}{\partial \theta_1} \mathcal{D}_1^n \frac{1}{f_\Lambda} \sum_{w \in S_N} \mathcal{K}_w \theta_{\{1, \dots, m\}} z^\Lambda. \quad (78)$$

We will now focus on the part of this expression involving $\theta_{\{1, \dots, m\}}$ to the left. This term is of the form

$$\mathcal{I}'_n m_\Lambda \Big|_{\theta_{\{1, \dots, m\}}} = \frac{1}{f_\Lambda} \sum_{w \in S_m} (-1)^{\ell(w)+1} \sum_{i=1}^m K_{1i} \mathcal{D}_1^n z^{w\Lambda}, \quad (79)$$

where $\ell(w)$ is the length of the permutation w . This is because, if $\kappa_w \theta_{\{1, \dots, m\}}$ does not contain θ_1 , the term will be annihilated by $\frac{\partial}{\partial \theta_1}$. Now, if $\kappa_w \theta_{\{1, \dots, m\}}$ contains θ_1 , then $\kappa_{1i} \kappa_w \theta_{\{1, \dots, m\}}$ contains θ_i , and thus $\kappa_{1i} \kappa_w \theta_{\{1, \dots, m\}}$ will certainly not be equal to $\pm \theta_{\{1, \dots, m\}}$ if $i > m$. Therefore $i \leq m$, which means that w needs to belong to S_m for $\kappa_{1i} \kappa_w \theta_{\{1, \dots, m\}}$ to be equal to $\pm \theta_{\{1, \dots, m\}}$. Finally, with $\kappa_{1i} \kappa_w \theta_{\{1, \dots, m\}} = (-1)^{\ell(w)+1} \theta_{\{1, \dots, m\}}$, formula (79) is seen to hold.

Now, if $w \in S_m$, then $z^{w\Lambda} \in L_\Lambda$, which implies, from Lemma 26, that $\mathcal{D}_i^n z^{w\Lambda} \in L_\Lambda$. Therefore, since K_{1i} , for $i = 1, \dots, m$, also preserves L_Λ (by definition), we have that $\mathcal{I}'_n m_\Lambda|_{\theta_{\{1, \dots, m\}}}$ belongs to L_Λ . Therefore, any z^Ω in $\mathcal{I}'_n m_\Lambda|_{\theta_{\{1, \dots, m\}}}$, for Ω a superpartition, will be such that $\Omega \leq \Lambda$, which proves the theorem. $\square$

Note that the eigenvalues $\epsilon_{\Lambda, n}$ are not given explicitly. However, $\epsilon_{\Lambda, 1}$ is obtained in the next theorem. This theorem also characterizes the precise action of $\mathcal{I}_1$ on monomials, if we discard coefficients that will not be needed in the sequel.

Theorem 28. *The action of $\mathcal{I}_1$ on the monomials is the following*

$$\mathcal{I}_1 m_\Lambda = \epsilon_{\Lambda, 1} m_\Lambda + \sum_{\Omega <_t \Lambda} b_{\Lambda\Omega} m_\Omega + \sum_{\Gamma <_h \Lambda} c_{\Lambda\Gamma} m_\Gamma, \quad (80)$$

with

$$\epsilon_{\Lambda,1} = (N-1)! \left[\sum_{i=1}^m \Lambda_i - \beta(m(m-1) + \#\Lambda) \right], \quad (81)$$

where $\#\Lambda$ is the number of pairs (i,j) such that $i \in \{1, \dots, m\}$, $j \in \{m+1, \dots, N\}$ and $\Lambda_i < \Lambda_j$, and with

$$b_{\Lambda\Omega} = \begin{cases} (N-1)! \beta \operatorname{sgn}(\sigma_{T_{ij}\Lambda}^a) n_{\Omega^s}(\Lambda_i) & \text{if } \Omega = \overline{T_{ij}\Lambda} \text{ for some } i < j \\ 0 & \text{otherwise} \end{cases} \quad (82)$$

(the $c_{\Lambda\Gamma}$'s are left undetermined).

The action of $\mathcal{I}_1$ given in (80) displays two types of subleading terms, each type being characterized by one of the two specializations of the Bruhat ordering. As in the action of $\mathcal{H}_n$, we recover terms that are h -ordered. But in addition, there appear terms that are t -ordered, labeled by superpartitions such that $\Lambda^* = \Omega^*$. It is precisely the superpolynomials associated to such superpartitions that were $\mathcal{H}_n$ -degenerate. It is because they can now be compared that the action of $\mathcal{I}_1$ lifts the degeneracies, as we will see in the lemma that follows this theorem.

Proof. To simplify the analysis, we again work with $\mathcal{I}'_1$ instead of $\mathcal{I}_1$, and focus on the coefficient $\theta_{\{1, \dots, m\}}$. From (79), this coefficient is given by

$$\mathcal{I}'_1 m_\Lambda \Big|_{\theta_{\{1, \dots, m\}}} = \frac{1}{f_\Lambda} \sum_{i=1}^m K_{1i} \mathcal{D}_1 \sum_{w \in S_m} (-1)^{\ell(w)+1} z^{w\Lambda}. \quad (83)$$

From the Hecke algebra relations, and $(1i) = \sigma_{i-1} \cdots \sigma_1 \cdots \sigma_{i-1}$ we obtain

$$K_{1i} \mathcal{D}_1 = \mathcal{D}_i K_{1i} + \beta \sum_{j=1}^{i-1} K_{(\sigma_{i-1} \cdots \sigma_1)_j \sigma_2 \cdots \sigma_{i-1}}, \quad (84)$$

where the symbol $()_j$ means that σ_j does not belong to the product in parenthesis. Now, the transposition $(1i)$ contains an odd number of elementary transpositions, while all the terms of the form $(\sigma_{i-1} \cdots \sigma_1)_j \sigma_2 \cdots \sigma_{i-1}$ contain an even number of such transpositions. With $\sum_{w \in S_m} (-1)^{\ell(w)+1} z^{w\Lambda}$ being totally antisymmetric in the first m variables, we thus obtain

$$\begin{aligned} \mathcal{I}'_1 m_\Lambda \Big|_{\theta_{\{1, \dots, m\}}} &= \frac{1}{f_\Lambda} \sum_{i=1}^m \left(\mathcal{D}_i - \beta(i-1) \right) \sum_{w \in S_m} (-1)^{\ell(w)} z^{w\Lambda} \\ &= \frac{1}{f_\Lambda} \sum_{w \in S_m} (-1)^{\ell(w)} K_w \sum_{i=1}^m \left(\mathcal{D}_i - \beta(i-1) \right) z^\Lambda, \end{aligned} \quad (85)$$

where we have used the fact that $\sum_{i=1}^m \mathcal{D}_i$ commutes with any K_w such that $w \in S_m$. We have, from (23),

$$\sum_{i=1}^m \left(\mathcal{D}_i - \beta(i-1) \right) = \sum_{i=1}^m z_i \frac{\partial}{\partial z_i} + \beta \sum_{1 \leq i < j \leq m} (\mathcal{O}_{ij} + \mathcal{O}_{ji}) + \beta \sum_{i=1}^m \sum_{j=m+1}^n \mathcal{O}_{ij} - \beta m(m-1). \quad (86)$$

With $\mathcal{O}_{ij} + \mathcal{O}_{ji} = 0$, this leads to

$$\mathcal{I}'_1 m_\Lambda \Big|_{\theta_{\{1, \dots, m\}}} = \left[\sum_{i=1}^m \Lambda_i - \beta m(m-1) \right] m_\Lambda + \frac{\beta}{f_\Lambda} \sum_{w \in S_m} (-1)^{\ell(w)} K_w \sum_{i=1}^m \sum_{j=m+1}^N \mathcal{O}_{ij} z^{w\Lambda}. \quad (87)$$

The last term of this expression becomes (if we do not consider coefficients that are not permutations of Λ)

$$\begin{aligned} \frac{\beta}{f_\Lambda} \sum_{w \in S_m} (-1)^{\ell(w)} K_w \sum_{i=1}^m \sum_{j=m+1}^N \mathcal{O}_{ij} z^\Lambda \\ = -\frac{\beta \#_\Lambda}{f_\Lambda} \sum_{w \in S_m} (-1)^{\ell(w)} K_w z^\Lambda + \frac{\beta}{f_\Lambda} \sum_{w \in S_m} (-1)^{\ell(w)} K_w \sum_{(i,j); \Lambda_i > \Lambda_j} z^{T_{ij}\Lambda}, \end{aligned} \quad (88)$$

where (i, j) is considered to be such that $i \in \{1, \dots, m\}$ and $j \in \{m+1, \dots, N\}$. Putting everything together, we get

$$\mathcal{I}'_1 m_\Lambda = \epsilon_{\Lambda,1} m_\Lambda + \frac{\beta}{f_\Lambda} \sum_{w \in S_N} K_w \theta_1 \cdots \theta_m \sum_{(i,j); \Lambda_i > \Lambda_j} z^{T_{ij}\Lambda} + \sum_{\Gamma <_h \Lambda} b_{\Lambda\Gamma} m_\Gamma. \quad (89)$$

If $\Omega = \overline{T_{ij}\Lambda}$, the coefficient of m_Ω in the last formula is then given by

$$\begin{aligned} \beta \operatorname{sgn}(\sigma_{T_{ij}\Lambda}^a) \frac{f_\Omega}{f_\Lambda} n_{\Lambda^s}(\Lambda_j) &= \beta \operatorname{sgn}(\sigma_{T_{ij}\Lambda}^a) \frac{n_{\Omega^s}(\Lambda_i)! n_{\Omega^s}(\Lambda_i)!}{n_{\Lambda^s}(\Lambda_j)! n_{\Lambda^s}(\Lambda_i)!} (n_{\Omega^s}(\Lambda_j) + 1) \\ &= \beta \operatorname{sgn}(\sigma_{T_{ij}\Lambda}^a) \frac{n_{\Omega^s}(\Lambda_i)}{n_{\Omega^s}(\Lambda_j) + 1} (n_{\Omega^s}(\Lambda_j) + 1) \\ &= \beta \operatorname{sgn}(\sigma_{T_{ij}\Lambda}^a) n_{\Omega^s}(\Lambda_i), \end{aligned} \quad (90)$$

since $n_{\Lambda^s}(\Lambda_i) = n_{\Omega^s}(\Lambda_i) - 1$ and $n_{\Lambda^s}(\Lambda_j) = n_{\Omega^s}(\Lambda_j) + 1$. $\square$

Lemma 29. *The triangular operator $\mathcal{I}_1$ is regular, that is, $\epsilon_{\Lambda,1} \neq \epsilon_{\Omega,1}$ if $\Omega < \Lambda$.*

Proof. $\Omega < \Lambda$ means that Ω differs from Λ by the application of a sequence of T_{ij} 's, with $i \leq m$ and $j > m$. It follows that $\sum_{i=1}^m \Omega_i < \sum_{i=1}^m \Lambda_i$, so that the constant term in $\epsilon_{\Omega,1}$, viewed as a polynomial in β , is strictly smaller than $\epsilon_{\Lambda,1}|_{\beta=0}$. This readily implies that $\epsilon_{\Omega,1} \neq \epsilon_{\Lambda,1}$. $\square$

Lemma 30. *The action of $\mathcal{I}_1$ on $\mathcal{J}_\Lambda$ is triangular with respect to the t -ordering, that is,*

$$\mathcal{I}_1 \mathcal{J}_\Lambda = \epsilon_{\Lambda,1} \mathcal{J}_\Lambda + \sum_{\Omega <_t \Lambda} b_{\Lambda\Omega} \mathcal{J}_\Omega, \quad (91)$$

where the coefficients $b_{\Lambda\Omega}$ are given in (82).

Proof. From Theorem 22 and because the order $\leq_h$ is weaker than the Bruhat order $\leq$, we have

$$\mathcal{J}_\Lambda = m_\Lambda + \sum_{\Omega < \Lambda; \Omega^* \neq \Lambda^*} v_{\Lambda\Omega} m_\Omega, \quad (92)$$

and thus, from (80),

$$\begin{aligned} \mathcal{I}_1 \mathcal{J}_\Lambda &= \epsilon_{\Lambda,1} m_\Lambda + \sum_{\Omega <_t \Lambda} b_{\Lambda\Omega} m_\Omega + \sum_{\Gamma < \Lambda; \Gamma^* \neq \Lambda^*} c_{\Lambda\Gamma} m_\Gamma \\ &= \epsilon_{\Lambda,1} \mathcal{J}_\Lambda + \sum_{\Omega <_t \Lambda} b_{\Lambda\Omega} \mathcal{J}_\Omega + \sum_{\Gamma < \Lambda; \Gamma^* \neq \Lambda^*} d_{\Lambda\Gamma} \mathcal{J}_\Gamma, \end{aligned} \quad (93)$$

since from (92) we get the inverse relation

$$m_\Lambda = \mathcal{J}_\Lambda + \sum_{\Omega < \Lambda; \Omega^* \neq \Lambda^*} w_{\Lambda\Omega} \mathcal{J}_\Omega. \quad (94)$$

It now suffices to show that the coefficients $d_{\Lambda\Gamma}$ in (93) do in fact vanish. Since $\mathcal{I}_1$ commutes with $\mathcal{H}_n$, $\mathcal{I}_1 \mathcal{J}_\Lambda$ must be an eigenfunction of $\mathcal{H}_n$ with eigenvalue $\varepsilon_{\Lambda,n}$ ($n = 1, 2, \dots$). From Theorem 23, the expansion of $\mathcal{I}_1 \mathcal{J}_\Lambda$ in terms of $\mathcal{J}_\Omega$ can thus only contain terms such that $\Omega^* = \Lambda^*$. $\square$

8 The orthogonal Jack superpolynomials J_Λ

Theorem 31. *There exists a unique basis $\{J_\Lambda\}_\Lambda$ of P^{S_N} such that*

$$\mathcal{I}_1 J_\Lambda = \epsilon_{\Lambda,1} J_\Lambda \quad \text{and} \quad J_\Lambda = \mathcal{J}_\Lambda + \sum_{\Omega <_t \Lambda} u_{\Lambda\Omega} \mathcal{J}_\Omega. \quad (95)$$

Proof. Using Lemmas 29 and 30, the theorem follows immediately from Theorem 16. $\square$

From Theorem 16, Lemma 30 and the explicit expression of the eigenvalues $\epsilon_{\Lambda,1}$ given in Theorem 28, a determinantal formula, giving J_Λ in terms of $\mathcal{J}_\Omega$, can be obtained. Moreover, by Corollary 17, a recurrence is provided for the coefficients $u_{\Lambda\Omega}$. Note that because the eigenvalues $\epsilon_{\Lambda,1}$ and the coefficients $b_{\Lambda,\Omega}$ given in Theorem 28 do not depend upon N , apart from a factorizable overall prefactor $(N-1)!$, the coefficients $u_{\Lambda\Omega}$ are N -independent. Moreover, since the expansion of $\mathcal{J}_\Lambda$ in the supermonomial basis is N -independent, this holds true for the supermonomial decomposition of J_Λ . This is illustrated in Appendix A.

Given Theorem 23, the previous theorem has the following corollary.

Corollary 32.

$$\mathcal{H}_n J_\Lambda = \varepsilon_{\Lambda,n} J_\Lambda. \quad (96)$$

Furthermore, if $\Lambda^* \neq \Omega^*$, then there exists at least one n such that $\varepsilon_{\Lambda,n} \neq \varepsilon_{\Omega,n}$.

We now show that J_Λ is also an eigenfunction of $\mathcal{I}_n$, for all n .

Theorem 33. *We have*

$$\mathcal{I}_n J_\Lambda = \epsilon_{\Lambda,n} J_\Lambda. \quad (97)$$

Furthermore, if $\Lambda^a \neq \Omega^a$, then there exists at least one n such that $\epsilon_{\Lambda,n} \neq \epsilon_{\Omega,n}$.

Proof. Given Theorem 31, formula (77) and the fact that $\mathcal{I}_n$ and $\mathcal{I}_1$ commute, the first part of the theorem follows immediately from Proposition 18. The second part of the theorem is obvious because the $\epsilon_{\Lambda,n}$'s ($n = 1, 2, \dots$) are polynomials in β whose constant terms $\epsilon_{\Lambda,n}|_{\beta=0} = \sum_{i=1}^m \Lambda_i^n$ are such that if $\epsilon_{\Lambda,n}|_{\beta=0} = \epsilon_{\Omega,n}|_{\beta=0}$ for $n = 1, 2, \dots$, then $\Lambda^a = \Omega^a$. Hence, $\epsilon_{\Lambda,n}$ and $\epsilon_{\Omega,n}$, considered as functions of a generic parameter β , cannot be equal when $\Lambda^a \neq \Omega^a$. $\square$

Now, it is obvious that if $\Lambda^* = \Omega^*$ and $\Lambda^a = \Omega^a$, then $\Lambda = \Omega$. Therefore, using Corollary 32, the previous theorem has the following corollary.

Corollary 34. *The polynomial J_Λ is the unique common eigenfunction of the operators $\mathcal{H}_n$ and $\mathcal{I}_\ell$ ($n, \ell = 1, 2, \dots$), with respective eigenvalues $\varepsilon_{\Lambda,n}$ and $\epsilon_{\Lambda,\ell}$.*

We thus have immediately, since the operators $\mathcal{H}_n$ and $\mathcal{I}_\ell$, $n, \ell = 1, 2, \dots$, are self-adjoint with respect to the scalar product $\langle \cdot, \cdot \rangle_\beta$, the orthogonality of the basis $\{J_\Lambda\}_\Lambda$.

Theorem 35. *The basis $\{J_\Lambda\}_\Lambda$ of P^{S_N} satisfies*

$$\langle J_\Lambda, J_\Omega \rangle_\beta = c_\Lambda(\beta) \delta_{\Lambda\Omega}, \quad (98)$$

where $c_\Lambda(\beta)$ is some function of β (to be determined in the next section).

9 Symmetrizing the non-symmetric Jack polynomials in super-space

For a composition λ , the non-symmetric Jack polynomials, E_λ , are the unique polynomials in the variables $z_1, \dots, z_N$ satisfying

$$E_\lambda = z^\lambda + \sum_{\mu < \lambda} c_{\lambda\mu}(\beta) z^\mu, \quad \text{and} \quad \langle E_\lambda, E_\mu \rangle_\beta \propto \delta_{\lambda\mu}. \quad (99)$$

The non-symmetric Jack polynomial E_λ is an eigenfunction of the Dunkl operators,

$$\mathcal{D}_i E_\lambda = \underline{\lambda}_i E_\lambda, \quad (100)$$

where the eigenvalue $\underline{\lambda}_i$ is given in (44). This property characterizes E_λ uniquely [6].

Using the orthogonality of the non-symmetric Jack polynomials, the polynomials $E_{\lambda,I} = \theta_I E_\lambda$ are immediately seen to form an orthogonal basis of P . These polynomials are in fact the unique common eigenfunctions of the operators $\mathcal{D}_i$ and $\theta_i \frac{\partial}{\partial \theta_i}$, for $i = 1, \dots, N$.

A basis of the space of S_N -symmetric polynomials in the variables $z_1, \dots, z_N$ is given by the Jack polynomials. The following formula is known [6, 7]

$$J_{\lambda^+} \propto \sum_{w \in S_N} K_w E_{\lambda^+}, \quad (101)$$

where J_{λ^+} is the Jack polynomial indexed by the partition λ^+ . On the operatorial side, the Jack polynomials are the unique common eigenfunctions of the operators $\mathcal{H}_n = \sum_{i=1}^N \mathcal{D}_i^n$ ($n = 1, 2, \dots$).

Other polynomials obtained from the non-symmetric Jack polynomials have been studied in [8, 9]. We are particularly interested in those obtained by antisymmetrizing the first m variables and symmetrizing the remaining ones. Namely, given partitions λ and μ with m and $N - m$ parts respectively, let⁸

$$S_{(\lambda,\mu)} = \frac{(-1)^{m(m-1)/2}}{f_\mu} \sum_{\sigma \in S_m} \sum_{w \in S_{m^c}} K_\sigma K_w (-1)^{\ell(\sigma)} E_{(\lambda^R, \mu^R)}, \quad (102)$$

where $E_{(\lambda^R, \mu^R)}$ is the non-symmetric Jack polynomial indexed by the concatenation of the compositions λ^R (recall that this is the partition λ in reversed order) and μ^R (that is, the adjunction of the entries of μ^R to the right of those of λ^R without rearrangement), and where S_{m^c} stands for the permutations of $\{m+1, \dots, N\}$ (or the permutations of S_N that leave $1, \dots, m$ fixed).

Property 36. [8]. *The polynomials $S_{(\lambda,\mu)}$ are orthogonal with respect to the scalar product $\langle \cdot, \cdot \rangle_\beta$, the norm being given explicitly by*

$$\langle S_{(\lambda,\mu)}, S_{(\lambda,\mu)} \rangle_\beta = \frac{m!(N-m)!}{f_\mu} \frac{d'_{(\lambda^R, \mu)} d_{(\lambda^R, \mu^R)}}{d'_{(\lambda^R, \mu^R)} d_{(\lambda, \mu^R)}} \langle E_{(\lambda^R, \mu^R)}, E_{(\lambda^R, \mu^R)} \rangle_\beta, \quad (103)$$

where for a composition γ (in this case given by the concatenation of two compositions)

$$\begin{aligned} d_\gamma &= \prod_{(i,j) \in \gamma} [a(i,j) + 1 + \beta(l(i,j) + 1)] \\ d'_\gamma &= \prod_{(i,j) \in \gamma} [a(i,j) + 1 + \beta l(i,j)] \end{aligned} \quad (104)$$

⁸In [8], the antisymmetrization (resp. symmetrization) is performed on the last m variables (resp. first $N - m$ variables). This does not affect in any meaningful way the properties of these polynomials. For instance, formula (103) can be extracted easily from a similar formula of [8].

with

$$\begin{aligned} a(i, j) &= \gamma_i - j \\ l(i, j) &= \#\{k = 1, \dots, i-1 \mid j \leq \gamma_k + 1 \leq \gamma_i\} + \#\{k = i+1, \dots, N \mid j \leq \gamma_k \leq \gamma_i\} \end{aligned} \quad (105)$$

and

$$\langle E_\gamma, E_\gamma \rangle_\beta = \prod_{1 \leq i < j}^N \prod_{p=0}^{\beta-1} \left(\frac{\gamma_j - \gamma_i + p}{\gamma_j - \gamma_i - p - 1} \right)^{\epsilon(\gamma_j - \gamma_i)}, \quad (106)$$

where $\epsilon(x) = 1$ if $x > 0$ and -1 otherwise.

We now build a basis of P^{S_N} from the non-symmetric Jack polynomials.

Definition 37. Given a superpartition $\Lambda = (\Lambda^a; \Lambda^s)$,

$$\tilde{J}_\Lambda = \frac{(-1)^{(m)(m-1)/2}}{f_{\Lambda^s}} \sum_{w \in S_N} \mathcal{K}_w \theta_{\{1, \dots, m\}} E_{((\Lambda^a)^R, (\Lambda^s)^R)}. \quad (107)$$

Proposition 38. We have

$$\tilde{J}_\Lambda = \sum_{w \in S_N / (S_m \times S_{m^c})} \mathcal{K}_w \theta_{\{1, \dots, m\}} S_{(\Lambda^a, \Lambda^s)}. \quad (108)$$

Proof: From the definition of $\tilde{J}_\Lambda$, we get

$$\begin{aligned} \tilde{J}_\Lambda &= \frac{(-1)^{(m)(m-1)/2}}{f_{\Lambda^s}} \sum_{w \in S_N / (S_m \times S_{m^c})} \mathcal{K}_w \sum_{\sigma \in S_m} \mathcal{K}_\sigma \sum_{\rho \in S_{m^c}} \mathcal{K}_\rho \theta_{\{1, \dots, m\}} E_{((\Lambda^a)^R, (\Lambda^s)^R)} \\ &= \sum_{w \in S_N / (S_m \times S_{m^c})} \mathcal{K}_w \theta_{\{1, \dots, m\}} \left(\frac{(-1)^{(m)(m-1)/2}}{f_{\Lambda^s}} \sum_{\substack{\sigma \in S_m \\ \rho \in S_{m^c}}} \mathcal{K}_\sigma \mathcal{K}_\rho (-1)^{\ell(\sigma)} E_{((\Lambda^a)^R, (\Lambda^s)^R)} \right), \end{aligned} \quad (109)$$

which gives the desired result from the definition of $S_{(\Lambda^a, \Lambda^s)}$. $\square$

Note that the left coset representatives of $S_N / (S_m \times S_{m^c})$ can be described as

$$\left\{ \sum_{k=0}^{\min(m, N-m)} \sum_{\substack{1 \leq i_1 < i_2 < \dots < i_k \leq m \\ m+1 \leq j_1 < j_2 < \dots < j_k \leq N}} \mathcal{K}_{i_1, j_1} \cdots \mathcal{K}_{i_k, j_k} \right\}, \quad (110)$$

with the understanding that when $k = 0$, the product of $\mathcal{K}$ factors reduces to the identity.

Proposition 39. We have

$$\tilde{J}_\Lambda = m_\Lambda + \sum_{\Omega < \Lambda} a_{\Lambda\Omega}(\beta) m_\Omega, \quad (111)$$

that is, $\tilde{J}_\Lambda$ is monic and triangularly related (with respect to the Bruhat ordering on superpartitions) to the monomial superfunction basis.

Proof: The monicity of $\tilde{J}_\Lambda$, given the monicity of $S_{(\lambda, \mu)}$ [8], follows from Proposition 38. Now, let Λ^R stand for $((\Lambda^a)^R, (\Lambda^s)^R)$. From Definition 37 and (99), the coefficient of $\theta_1 \cdots \theta_m$ in $\tilde{J}_\Lambda$ is

$$\begin{aligned} \tilde{J}_\Lambda \Big|_{\theta_1 \cdots \theta_m} &= \frac{(-1)^{m(m-1)/2}}{f_{\Lambda^s}} \sum_{\sigma \in S_m} \sum_{w \in S_{m^c}} \mathcal{K}_\sigma \mathcal{K}_w (-1)^{\ell(\sigma)} E_{\Lambda^R} \\ &= \frac{(-1)^{m(m-1)/2}}{f_{\Lambda^s}} \sum_{\sigma \in S_m} \sum_{w \in S_{m^c}} \mathcal{K}_\sigma \mathcal{K}_w (-1)^{\ell(\sigma)} \sum_{\Omega \leq \Lambda^R} c_{\Lambda\Omega}(\beta) z^\Omega, \end{aligned} \quad (112)$$

where in the last equality, the ordering is on compositions. To obtain the monomial superfunctions that appear in the expansion of $\tilde{J}_\Lambda$, we must simply select the superpartitions that arise as powers of z in this last equality. Because of the nature of the sums over σ and w , the only superpartition Γ such that $\Gamma^* = \Lambda^*$ that can arise is Λ . Finally, since any rearrangement Γ of a composition Ω such that $\Omega < \Lambda^R$ and $\Omega^* \neq \Lambda^*$ is also such that $\Gamma < \Lambda$, the proposition is seen to hold. $\square$

The orthogonality of the $\tilde{J}_\Lambda$'s is almost immediate from Proposition 38.

Proposition 40. *We have*

$$\langle \tilde{J}_\Lambda, \tilde{J}_\Omega \rangle_\beta = \delta_{\Lambda\Omega} \frac{N!}{m!(N-m)!} \langle S_{(\Lambda^a, \Lambda^s)}, S_{(\Lambda^a, \Lambda^s)} \rangle_\beta, \quad (113)$$

where $\langle S_{(\Lambda^a, \Lambda^s)}, S_{(\Lambda^a, \Lambda^s)} \rangle_\beta$ is given explicitly in (103).

Proof: Using Proposition 38, we have

$$\langle \tilde{J}_\Lambda, \tilde{J}_\Omega \rangle_\beta = \left\langle \sum_{w \in G} \mathcal{K}_w \theta_{\{1, \dots, m\}} S_{(\Lambda^a, \Lambda^s)}, \sum_{\sigma \in G} \mathcal{K}_\sigma \theta_{\{1, \dots, m\}} S_{(\Omega^a, \Omega^s)} \right\rangle_\beta, \quad (114)$$

where G is a set of left coset representatives of $S_N / (S_m \times S_{m^c})$. Since $\mathcal{K}_\sigma^\dagger = \mathcal{K}_{\sigma^{-1}}$, this gives

$$\langle \tilde{J}_\Lambda, \tilde{J}_\Omega \rangle_\beta = \left\langle \sum_{w, \sigma \in G} \mathcal{K}_{\sigma^{-1}} \mathcal{K}_w \theta_{\{1, \dots, m\}} S_{(\Lambda^a, \Lambda^s)}, \theta_{\{1, \dots, m\}} S_{(\Omega^a, \Omega^s)} \right\rangle_\beta. \quad (115)$$

Now, $\mathcal{K}_{\sigma^{-1}} \mathcal{K}_w$ must belong to $S_m \times S_{m^c}$ for the product of θ 's to be the same on both sides. Therefore, since this only occurs for $w = \sigma$,

$$\langle \tilde{J}_\Lambda, \tilde{J}_\Omega \rangle_\beta = \#G \langle S_{(\Lambda^a, \Lambda^s)}, S_{(\Omega^a, \Omega^s)} \rangle_\beta, \quad (116)$$

which proves the proposition since the cardinality of G is $N! / (m!(N-m)!)$. $\square$

We now make the connection with the family $\{J_\Lambda\}_\Lambda$ introduced before.

Theorem 41. *For any superpartition Λ , we have $J_\Lambda = \tilde{J}_\Lambda$.*

Proof: Both families are triangular with respect to the same ordering (namely, the Bruhat ordering on compositions) when expanded in the supermonomial basis. Since the Gram-Schmidt orthonormalization procedure ensures that there exists at most one orthonormal family with such triangularity, the two families can only differ by a constant. The theorem then follows from the monicity of both families. $\square$

10 Conclusion

This work has many natural generalizations. The most direct one is to consider the rational counterpart of the above results. The orthogonal eigenfunctions of the supersymmetric rational CMS model can be obtained in a rather direct way using the present results and the remarkable relation, preserved in the supersymmetric case, that exists between the eigenfunctions of the trigonometric and rational models [13]. This leads to a closed form expression (a determinant of determinants) for the orthogonal generalized Hermite (or Hi-Jack) superpolynomials. These results will be presented in [14].

Another rather immediate line of generalizations would be to examine the supersymmetric extension of the r/tCMS models associated to root systems of any type. To find the corresponding orthogonal superpolynomials, one would proceed as follows: take the Dunkl operator of the corresponding exchange version of the r/tCMS model of interest, look for their non-symmetric eigenfunctions, dress them with a fermionic monomial prefactor and symmetrize the result with respect to both types of variables. For instance, the generalized Jacobi and Laguerre polynomials in superspace could be constructed in this way. The resulting superpolynomial would be a linear combination (with θ_I coefficients) of the corresponding version (Jacobi or Laguerre) of the generalized polynomials with mixed symmetry [9]. Note, however, that the norm of these special polynomials has not yet been computed. The conserved charges of the model would be constructed exactly as in the present case, by symmetrizing the Dunkl operators raised to the n -th power and multiplied by a fermionic prefactor.

Along these lines, the formulation of the supersymmetric extension of the elliptic (eCMS) model appears to be direct. The elliptic Dunkl operators are given in [15]. Their supersymmetric lift is immediate, leading directly to an expression for the Hamiltonian of the seCMS model. Again, the general form of the charges is bound to be similar to the one found in this article. However, in this case, little is known about the eigenfunctions.

The q deformation of the Jack polynomials are the Macdonald polynomials [3], eigenfunctions (up to a conjugation) of the Ruijsenaars-Schneider model [16], a relativistic version of the CMS model. Again, there exist q analogues of the Dunkl operators, giving a natural road for the formulation of the supersymmetric Ruijsenaars-Schneider model. Moreover, the nonsymmetric Macdonald polynomials being known, they can be lifted to orthogonal Macdonald superpolynomials.

Further avenues regarding future works concern the study of properties of the Jack superpolynomials J_Λ themselves. On that matter, we already have strong indications that these objects have rather remarkable properties. In particular, the Pieri formulas appear to be rather nice. Moreover, exploratory analyses indicate that J_Λ products have a combinatorial interpretation in terms of novel types of supertableaux. Finally, a natural problem that should not be out of reach at this stage is working out the superspace extension of the operator construction of [1].

ACKNOWLEDGMENTS

This work was supported by NSERC. L.L. wishes to thank Luc Vinet for his support, and P.D. is grateful to the Fondation J.A Vincent for a student fellowship.

A Examples of Jack superpolynomials

In this appendix, we present a detailed calculation of one Jack polynomial in superspace via determinantal formulas. Then, we give explicitly all orthogonal superpolynomials of degrees not larger than 3.

Let $\Lambda = (3, 1; 0)$. It has weight 4 and lies in the 2-fermion sector. The explicit action of the conserved operator $\mathcal{I}_1$ in the space $\text{Span}\{\mathcal{J}_\Omega\}_{\Omega <_t \Lambda}$ is obtained from Lemma 30:

$$\begin{aligned}\mathcal{I}_1 \mathcal{J}_{(3,1;0)} &= (N-1)!(4-2\beta) \mathcal{J}_{(3,1;0)} + (N-1)! \beta \mathcal{J}_{(3,0;1)} - (N-1)! \beta \mathcal{J}_{(1,0;3)}, \\ \mathcal{I}_1 \mathcal{J}_{(3,0;1)} &= (N-1)!(3-3\beta) \mathcal{J}_{(3,0;1)} + (N-1)! \beta \mathcal{J}_{(1,0;3)},\end{aligned}$$

$$\mathcal{I}_1 \mathcal{J}_{(1,0;3)} = (N-1)!(1-4\beta) \mathcal{J}_{(1,0;3)}. \quad (117)$$

Theorem 16 allows to express the orthogonal superpolynomial $J_{(3,1;0)}$ as the following determinant:

$$\begin{aligned} J_{(3,1;0)} &= \frac{1}{(N-1)!(3+2\beta)(N-1)!(1+\beta)} \begin{vmatrix} \mathcal{J}_{(1,0;3)} & -(N-1)!(3+2\beta) & 0 \\ \mathcal{J}_{(3,0;1)} & (N-1)!\beta & -(N-1)!(1+\beta) \\ \mathcal{J}_{(3,1;0)} & -(N-1)!\beta & (N-1)!\beta \end{vmatrix} \\ &= \mathcal{J}_{(3,1;0)} + \frac{\beta}{1+\beta} \mathcal{J}_{(3,0;1)} - \frac{\beta}{(3+2\beta)(1+\beta)} \mathcal{J}_{(1,0;3)}. \end{aligned} \quad (118)$$

We now want the monomial decomposition of the previous result. For this, we must determine the action of the Hamiltonian $\mathcal{H}_2$ in the space $\text{Span}\{m_\Omega\}_{\Omega <_h \Lambda}$. Using Theorem 2 in Ref. [5], the action of $\mathcal{H}_2$ on the monomial $m_{(3,1;0)}$ is given by

$$\mathcal{H}_2 m_{(3,1;0)} = (10 - 6\beta + 4N\beta) m_{(3,1;0)} + 4\beta m_{(2,1;1)}, \quad (119)$$

The monomial $m_{(2,1;1)}$ is itself an eigenfunction of the Hamiltonian, i.e.,

$$\mathcal{H}_2 m_{(2,1;1)} = (6 - 10\beta + 4N\beta) m_{(2,1;1)}. \quad (120)$$

Again, Theorem 16 yields the non-orthogonal superpolynomial $\mathcal{J}_{(3,1;0)}$ as a determinant:

$$\begin{aligned} \mathcal{J}_{(3,1;0)} &= \frac{1}{(4\beta)(-4-4\beta)} \begin{vmatrix} m_{(2,1;1)} & -4-4\beta \\ m_{(2,1;0)} & 4\beta \end{vmatrix} \\ &= m_{(3,1;0)} + \frac{\beta}{1+\beta} m_{(2,1;1)} \end{aligned} \quad (121)$$

Proceeding in the same way, we get:

$$\begin{aligned} \mathcal{J}_{(3,0;1)} &= \frac{1}{(-6-10\beta)(-4-4\beta)^3(-2-2\beta)} \begin{vmatrix} m_{(1,0;1^3)} & -6-10\beta & 0 & 0 & 0 & 0 \\ m_{(1,0;2,1)} & 12\beta & -4-4\beta & 0 & 0 & 0 \\ m_{(2,0;1^2)} & 6\beta & 0 & -4-4\beta & 0 & 0 \\ m_{(2,1;1)} & 0 & 0 & 0 & -4-4\beta & 0 \\ m_{(2,0;2)} & 0 & 2\beta & 4\beta & 2\beta & -2-2\beta \\ m_{(3,0;1)} & 0 & 2\beta & 8\beta & 2\beta & 2\beta \end{vmatrix} \\ &= m_{(3,0;1)} + \frac{\beta}{1+\beta} m_{(2,0;2)} + \frac{\beta(1+2\beta)}{2(1+\beta)^2} m_{(2,1;1)} \\ &\quad + \frac{\beta(2+3\beta)}{(1+\beta)^2} m_{(2,0;1^2)} + \frac{\beta(1+2\beta)}{2(1+\beta)^2} m_{(1,0;2,1)} + \frac{3\beta^2}{(1+\beta)^2} m_{(1,0;1^3)} \end{aligned} \quad (122)$$

and

$$\begin{aligned} \mathcal{J}_{(1,0;3)} &= \frac{1}{(-6-10\beta)(-4-4\beta)^3(-2-2\beta)} \begin{vmatrix} m_{(1,0;1^3)} & -6-10\beta & 0 & 0 & 0 & 0 \\ m_{(1,0;2,1)} & 12\beta & -4-4\beta & 0 & 0 & 0 \\ m_{(2,0;1^2)} & 6\beta & 0 & -4-4\beta & 0 & 0 \\ m_{(2,1;1)} & 0 & 0 & 0 & -4-4\beta & 0 \\ m_{(2,0;2)} & 0 & 2\beta & 4\beta & 2\beta & -2-2\beta \\ m_{(1,0;3)} & 0 & 6\beta & 0 & -2\beta & 2\beta \end{vmatrix} \\ &= m_{(3,0;1)} + \frac{\beta}{1+\beta} m_{(2,0;2)} - \frac{\beta}{2(1+\beta)^2} m_{(2,1;1)} \\ &\quad + \frac{\beta^2}{(1+\beta)^2} m_{(2,0;1^2)} + \frac{\beta(3+4\beta)}{2(1+\beta)^2} m_{(1,0;2,1)} + \frac{3\beta^2}{(1+\beta)^2} m_{(1,0;1^3)}. \end{aligned} \quad (123)$$

Consequently, the orthogonal Jack superpolynomial $J_{(3,1;0)}$ is written in the monomial basis as:

$$\begin{aligned}
J_{(3,1;0)} = & m_{(3,1;0)} + \frac{\beta}{1+\beta} m_{(3,0;1)} - \frac{\beta}{(1+\beta)(3+2\beta)} m_{(1,0;3)} \\
& + \frac{2\beta^2}{(1+\beta)(3+2\beta)} m_{(2,0;2)} + \frac{\beta(3+4\beta)}{(1+\beta)(3+2\beta)} m_{(2,1;1)} \\
& + \frac{6\beta^2}{(1+\beta)(3+2\beta)} m_{(2,0;1^2)} + \frac{2\beta^3}{(1+\beta)^2(3+2\beta)} m_{(1,0;2,1)} + \frac{6\beta^3}{(1+\beta)^2(3+2\beta)} m_{(1,0;1^3)}. \quad (124)
\end{aligned}$$

Finally, we give the simplest Jack superpolynomials explicitly in Tables 1 and 2. The polynomials have degrees less than 3 and 4 in θ and z respectively. In the first table, the non-orthogonal eigenfunctions $\mathcal{J}_\Lambda$ are written in the monomial basis $\{m_\Omega\}_{\Omega <_h \Lambda}$. The second table presents the orthogonal Jack polynomials in the non-orthogonal basis $\{\mathcal{J}_\Omega\}_{\Omega <_t \Lambda}$. We stress that these expressions do not depend on the number of variables N .

Table 1: Non-orthogonal superpolynomials $\mathcal{J}_\Lambda$ of weight $|\Lambda| \leq 3$

Superpartition Λ	Superpolynomial $\mathcal{J}_\Lambda(z, \theta; 1/\beta)$
(0)	$m_{(0)}$
(1)	$m_{(1)}$
(1 ²)	$m_{(1^2)}$
(2)	$m_{(2)} + \frac{2\beta}{1+\beta}m_{(1^2)}$
(1 ³)	$m_{(1^3)}$
(2, 1)	$m_{(2,1)} + \frac{6\beta}{1+2\beta}m_{(1^3)}$
(3)	$m_{(3)} + \frac{3\beta}{2+\beta}m_{(2,1)} + \frac{6\beta^2}{(1+\beta)(2+\beta)}m_{(1^3)}$
(0; 0)	$m_{(0;0)}$
(0; 1)	$m_{(0;1)}$
(1; 0)	$m_{(1;0)}$
(0; 1 ²)	$m_{(0;1^2)}$
(1; 1)	$m_{(1;1)}$
(0; 2)	$m_{(0;2)} + \frac{\beta}{1+\beta}m_{(1;1)} + \frac{2\beta}{1+\beta}m_{(0;1^2)}$
(2; 0)	$m_{(2;0)} + \frac{\beta}{1+\beta}m_{(1;1)}$
(0; 1 ³)	$m_{(0;1^3)}$
(1; 1 ²)	$m_{(1;1^2)}$
(0; 2, 1)	$m_{(0;2,1)} + \frac{2\beta}{1+2\beta}m_{(1;1^2)} + \frac{6\beta}{1+2\beta}m_{(0;1^3)}$
(1; 2)	$m_{(1;2)} + \frac{2\beta}{1+2\beta}m_{(1;1^2)}$
(2; 1)	$m_{(2;1)} + \frac{2\beta}{1+2\beta}m_{(1;1^2)}$
(0; 3)	$m_{(0;3)} + \frac{\beta}{2+\beta}m_{(2;1)} + \frac{2\beta}{2+\beta}m_{(1;2)} + \frac{3\beta}{2+\beta}m_{(0;2,1)} + \frac{4\beta^2}{(1+\beta)(2+\beta)}m_{(1;1^2)} + \frac{6\beta^2}{(1+\beta)(2+\beta)}m_{(0;1^3)}$
(3; 0)	$m_{(3;0)} + \frac{2\beta}{2+\beta}m_{(2;1)} + \frac{\beta}{2+\beta}m_{(1;2)} + \frac{2\beta^2}{(1+\beta)(2+\beta)}m_{(1;1^2)}$
(1, 0; 0)	$m_{(1,0;0)}$
(1, 0; 1)	$m_{(1,0;1)}$
(2, 0; 0)	$m_{(2,0;0)} + \frac{\beta}{1+\beta}m_{(1,0;1)}$
(1, 0; 1 ²)	$m_{(1,0;1^2)}$
(1, 0; 2)	$m_{(1,0;2)} + \frac{2\beta}{1+2\beta}m_{(1,0;1^2)}$
(2, 0; 1)	$m_{(2,0;1)} + \frac{2\beta}{1+2\beta}m_{(1,0;1^2)}$
(2, 1; 0)	$m_{(2,1;0)}$
(3, 0; 0)	$m_{(3,0;0)} + \frac{\beta}{2+\beta}m_{(2,1;0)} + \frac{2\beta}{2+\beta}m_{(2,0;1)} + \frac{\beta}{2+\beta}m_{(1,0;2)} + \frac{2\beta^2}{(1+\beta)(2+\beta)}m_{(1,0;1^2)}$

Table 2: Orthogonal Jack superpolynomials J_Λ of weight $|\Lambda| \leq 3$

Superpartition Λ	Jack superpolynomial $J_\Lambda(z, \theta; 1/\beta)$
(0)	$\mathcal{J}_{(0)}$
(1)	$\mathcal{J}_{(1)}$
(1 ²)	$\mathcal{J}_{(1^2)}$
(2)	$\mathcal{J}_{(2)}$
(1 ³)	$\mathcal{J}_{(1^3)}$
(2, 1)	$\mathcal{J}_{(2,1)}$
(3)	$\mathcal{J}_{(3)}$
(0; 0)	$\mathcal{J}_{(0;0)}$
(0; 1)	$\mathcal{J}_{(0;1)}$
(1; 0)	$\mathcal{J}_{(1;0)} + \frac{\beta}{1+\beta} \mathcal{J}_{(0;1)}$
(0; 1 ²)	$\mathcal{J}_{(0;1^2)}$
(1; 1)	$\mathcal{J}_{(1;1)} + \frac{2\beta}{1+2\beta} \mathcal{J}_{(0;1^2)}$
(0; 2)	$\mathcal{J}_{(0;2)}$
(2; 0)	$\mathcal{J}_{(2;0)} + \frac{\beta}{2+\beta} \mathcal{J}_{(0;2)}$
(0; 1 ³)	$\mathcal{J}_{(0;1^3)}$
(1; 1 ²)	$\mathcal{J}_{(1;1^2)} + \frac{3\beta}{1+3\beta} \mathcal{J}_{(0;1^3)}$
(0; 2, 1)	$\mathcal{J}_{(0;2,1)}$
(1; 2)	$\mathcal{J}_{(1;2)} + \frac{\beta}{1+\beta} \mathcal{J}_{(0;2,1)}$
(2; 1)	$\mathcal{J}_{(2;1)} + \frac{\beta}{1+\beta} \mathcal{J}_{(1;2)} + \frac{\beta(1+2\beta)}{2(1+\beta)^2} \mathcal{J}_{(0;2,1)}$
(0; 3)	$\mathcal{J}_{(0;3)}$
(3; 0)	$\mathcal{J}_{(3;0)} + \frac{\beta}{3+\beta} \mathcal{J}_{(0;3)}$
(1, 0; 0)	$\mathcal{J}_{(1,0;0)}$
(1, 0; 1)	$\mathcal{J}_{(1,0;1)}$
(2, 0; 0)	$\mathcal{J}_{(2,0;0)}$
(1, 0; 1 ²)	$\mathcal{J}_{(1,0;1^2)}$
(1, 0; 2)	$\mathcal{J}_{(1,0;2)}$
(2, 0; 1)	$\mathcal{J}_{(2,0;1)} + \frac{\beta}{1+\beta} \mathcal{J}_{(1,0;2)}$
(2, 1; 0)	$\mathcal{J}_{(2,1;0)} + \frac{\beta}{1+\beta} \mathcal{J}_{(1,0;2)} - \frac{\beta}{2(1+\beta)^2} \mathcal{J}_{(1,0;2)}$
(3, 0; 0)	$\mathcal{J}_{(3,0;0)}$

References

- [1] L. Lapointe and L. Vinet, *Exact operator solution of the Calogero-Sutherland model*, Commun. Math. Phys. **178**, 425 (1996); q-alg/9509003.
- [2] R. P. Stanley, *Some combinatorial properties of Jack symmetric functions*, Adv. Math. **77**, 76 (1988).
- [3] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., The Clarendon Press Oxford University Press, (1995).
- [4] P. Desrosiers, L. Lapointe and P. Mathieu, *Supersymmetric Calogero-Moser-Sutherland models and Jack superpolynomials*, Nucl. Phys. **B606**, 547 (2001); hep-th/0103178.
- [5] P. Desrosiers, L. Lapointe and P. Mathieu, *Jack superpolynomials, superpartition ordering and determinantal formulas*, Commun. Math. Phys. **233**, 383 (2003); hep-th/0105107.
- [6] E. Opdam, *Harmonic analysis for certain representations of graded hecke algebras*, Acta Math. **175**, 75 (1995).
- [7] F. Knop and S. Sahi, *A recursion and a combinatorial formula for the Jack polynomials*, Invent. Math. **128**, 9 (1997).
- [8] T. H. Baker, C. F. Dunkl and P. J. Forrester, *Polynomial eigenfunctions of the Calogero-Sutherland model with exchange terms*, in *Calogero-Moser-Sutherland Models*, ed. by J. F. van Diejen and L. Vinet, Springer (2000), 37.
- [9] T. H. Baker, and P. J. Forrester, *The Calogero-Sutherland model and polynomial with prescribed symmetry*, Nucl. Phys. **B492**, 682 (1997).
- [10] I. Cherednik, *A unification of Knizhnik-Zamolodchikov equation and Dunkl operators via affine Hecke algebras*, Inven. Math. **106**, 411 (1991).
- [11] J.-F. van Diejen, L. Lapointe, and J. Morse, *Determinantal construction of orthogonal polynomials associated with root systems*, submitted, math.CO/0303263.
- [12] L. Lapointe, A. Lascoux and J. Morse, *Determinantal formula and recursion for Jack polynomials*, Electro. J. Comb. **7**, 467 (2000).
- [13] M. Lasalle, *Polynômes de Hermite généralisés*, C.R. Acad. Sci. Paris, t. 313, séries I (1991), 579; K. Sogo, *A simple derivation of multivariable Hermite and Legendre polynomials*, J. Phys. Soc. Jap. **65**, 3097 (1996).
- [14] P. Desrosiers, L. Lapointe and P. Mathieu, *Generalized Hermite polynomials in superspace as eigenfunctions of the supersymmetric rational CMS model*, to appear in Nucl. Phys. B; hep-th/0305038; *Explicit formulas for the generalized Hermite polynomials in superspace*, hep-th/0309067.
- [15] V. M. Buchstaber, G. Felder, A. V. Veselov, *Elliptic Dunkl operators, root systems, and functional equations*, Duke Math. J. **76**, 885 (1994).
- [16] S. N. Ruijsenaars and H. Schneider *A new class of integrable systems and its relation to solitons*, Ann.Phys. **170**, 370 (1986) ; S. N. Ruijsenaars, *Complete integrability of relativistic Calogero-Moser systems and elliptic function identities*, Commun. Math. Phys. **110**, 191 (1987).